

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re patent application of Glennon and Hellburg

Serial No. 10/526,076

Group Art Unit 1625

Conf. #: 8312

Filed: October 24, 2005

Examiner Covington

For "β-HYDROXYPHENYLALKYLAMINES AND THEIR USE FOR TREATING
GLAUCOMA"

Commissioner for Patents

P.O. Box 1450

Alexandria, VA 22313-1450

DECLARATION UNDER 37 C.F.R. § 1.132
OF DR. RICHARD A. GLENNON

Sir:

1. I hold a PhD in Medicinal Chemistry (1973) from the School of Pharmacy at the State University of New York in Buffalo, New York.

2. As can be seen from my attached *Curriculum vitae*, I have held a number of positions in the field of medicinal chemistry throughout my career, and I am currently Chair of the Department of Medicinal Chemistry at Virginia Commonwealth University in Richmond, Virginia. I am considered an international expert in drug design. I have authored more than 350 peer-reviewed articles and more than 50 textbook chapters. I have edited several books related to medicinal chemistry, and I am inventor of 13 issued United States patents. I thus qualify as an "expert" in the field of medicinal chemistry and drug design, and I am able to provide evidence on matters pertaining to medicinal chemistry and on the level of one of ordinary skill in the art.

3. I am an inventor of and have reviewed the subject patent application, including the claims, and the Examiner's remarks as contained in the Office Action mailed on January 22, 2009.

4. Regarding the Examiner's opinion that substitutions in a molecule, such as replacement of a hydrogen atom with methyl group, is normally within the sphere of obviousness that surrounds a known compound, it is my expert opinion that while this may sometimes be the case, there are many instances where this is not true and thus this conclusion should not be

adopted as a general rule. In addition, it is my opinion that one of ordinary skill in the art, being a person with a doctoral degree, 5-10 years of research experience in medicinal chemistry or related fields, and an author of 10 or more peer-reviewed articles, would recognize that in many cases, replacement of a hydrogen atom with a methyl group can have a dramatic effect on the performance of a molecule. For example, the attached article (Glennon et al., J. Med. Chem, 2000. 43. 1011-1018, copy enclosed) teaches that known 5HT₃ receptors do not readily accommodate a tryptamine 5-methoxy group. For example, 5-methoxy tryptamine (compound 1b in the article), the O-methyl ether of 5-HT (compound 1a in the article, also known as serotonin), is completely devoid of activity at 5-HT₃ receptors whereas 1a (serotonin) readily binds to 5-HT₃ receptors (see the first paragraph of the Results and Discussion). In other words, the presence or absence of the methyl group, versus H, makes a large difference in the activity of the compound, and this difference would not be obvious to one of skill in the art merely from an examination of the chemical formulas. The differences were observed only when experimental results were obtained. In addition, data obtained in the experiments described in Glennon et al., show other examples of large impacts on activity when changes such as H to methyl are carried out. Compounds 11 and 12 in the table on page 1013 display 5-HT₆ receptor affinities (K_i's) of 78 vs 510, respectively. The difference between 11 and 12 is that 11 has H at position R1 whereas the R1 position is methyl in 12. In other words, we have conclusively demonstrated that H is NOT equal to Me in all instances! In my opinion, this data demonstrates and would demonstrate to one of ordinary skill in the art, that small changes in variable groups in a molecule, such as the substitution of H with methyl, can have a significant effect on the activity of a molecule.


5. With specific regards to the present invention, it would not be obvious to one of skill in the art, with a knowledge of Furukawa, Shell, Chiou and Bodor, to use the compounds depicted in Formula I of the present application to treat glaucoma. The compounds of Furukawa differ from those of Shell, Chiou and Bodor and those of the present invention (at least) by not having a methyl group on the ethylamine chain, and Furukawa does not teach the treatment of glaucoma but the treatment of obesity and/or diabetes. The compounds of Shell, Chiou and Bodor differ from those used in the methods of the present invention (at least) by having H *para* to the ethylamine chain, rather than C₁₋₃ alkyl, Cl, Br, or I, as is required in the practice of the present invention. Due to the sensitivity of biological systems, where even small changes in the structure

of a compound can have unpredictable results, one of skill in the art with a knowledge of Furukawa, Shell, Chiou and Bodor would not assume that the compounds used in the methods of the present invention would be useful to treat glaucoma.

6. In addition, with specific regard to the present invention, the experimental results demonstrate that the presence of a methyl on the ethylamine side chain of the compounds utilized in the present invention confers stability to the compounds. These results show that the presence of the methyl group instead of H confers significant metabolic stability on the molecule, which is otherwise susceptible to oxidative deamination by MAO. This provides another example of how a change from H to methyl can have far-reaching, non-obvious effects on the properties of a molecule.

7. I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the above-referenced application and any patent issuing thereon.

Date 4/24/09

Signed 
Dr. Richard A. Glennon

RICHARD A. GLENNON

PERSONAL INFORMATION

Home Address: 3810 Reeds Landing Circle
Midlothian, Virginia 23113

Office Address: **554A Smith Building; Department of Medicinal Chemistry
School of Pharmacy, Box 980540; Medical College of Virginia Campus
Virginia Commonwealth University, Richmond, VA 23298-0540**

Phone: (804)828-8487
Fax: (804)828-7404
E-mail: glennon@vcu.edu

EDUCATION

Ph.D. 1973: Medicinal Chemistry; School of Pharmacy,
State University of New York; Buffalo, NY

M. S. 1969: Medicinal Chemistry; School of Pharmacy
Northeastern University; Boston, MA

B. S. 1967: Pharmacy; Northeastern University; Boston, MA

POSTDOCTORAL TRAINING

Alcohol, Drug Abuse, Mental Health Administration (ADAMHA) Postdoctoral Fellow (Psychopharmacology);
Department of Pharmacology and Experimental Therapeutics, School of Medicine; State University of New
York: 1973 - 1975.

PROFESSIONAL POSITIONS

- Chair, Department of Medicinal Chemistry, VCU, 2006- to date
- Interim Chair, Department of Medicinal Chemistry, VCU, 2005- 2006
- Vice Chair, Department of Medicinal Chemistry, VCU, 1998-2005
- Acting Chair, Department of Medicinal Chemistry, MCV/VCU, 1997
- Associate Dept. Chair, Department of Medicinal Chemistry, School of Pharmacy, VCU; 1990-1996
- Graduate Program Director; Department of Medicinal Chemistry, VCU, 1995- 2007
- Professor, Department of Medicinal Chemistry School of Pharmacy, VCU; 1983- to date
- Affiliate Professor, Department of Pharmacology and Toxicology, School of Medicine; VCU; 1986- to date
- Associate Professor, Department of Medicinal Chemistry, School of Pharmacy, VCU; 1980-1983
- Assistant Professor, Department of Medicinal Chemistry, School of Pharmacy, VCU; 1975-1980
- Research Chemist, Warner-Lambert Research Institute; Morris Plains New Jersey; 1967-1968
- Staff Pharmacist, Bon Secours Hospital, Methuen, MA 1969; Staff Pharmacist, Childrens' Hospital and Medical Center, Boston, Massachusetts 1967 (Pharmacy Intern 1962-1966); Community Pharmacist, Part-time: Boston, MA; Lawrence, MA; Methuen, MA; Brookline, MA. 1967-1969

HONORS

- School of Pharmacy *Instructor of the Year Award*, 1979
- Virginia Commonwealth University *Distinguished Scholar Award*, 1993
- American Pharmaceutical Association *Research Achievement Award*, 1995
- Virginia Commonwealth University *Laboratory Safety Award* – 2000, 2004, 2006
- Florida A&M University, *Center of Excellence Distinguished Lecturer Award* – 2003
- Virginia Commonwealth University, School of Pharmacy *Instructor of the Year Award*, 2004
- Virginia Commonwealth University, School of Pharmacy *Research Award*, 2005
- Virginia Commonwealth University, *Award of Excellence*, 2004
- European *Order of the Oak and Tulip* award for excellence in receptor medicinal chemistry, 2007

PUBLICATIONS

More than 400 scientific publications and book chapters

EDITORIAL BOARD ACTIVITIES

- Senior Editor, *Journal of Medicinal Chemistry*
- Member, Editorial Board of *Pharmacology, Biochemistry & Behavior*
- Member, Editorial Board of *Current Medicinal Chemistry: CNS Agents*
- Member, Editorial Board *Current Topics in Medicinal Chemistry*
- Editor Emeritus and Consultant for *Medicinal Chemistry Research*
- Member, Editorial Board of *Brazilian Journal of Pharmaceutical Sciences*
- Associate Editor of *Journal of Drug Education and Awareness*
- Member, Editorial Board, *Current Chemical Biology*
- Member, Editorial Board, *Medicinal Chemistry Reviews*
- Member, Editorial Board, *Burger's Medicinal Chemistry*

PUBLICATIONS

1973-1979

1. Coburn, R. A.; Carapellotti, R. A.; Glennon, R. A. A PPP π -SCF variable integral study of mesoionic analogs based on six-membered ring mesoionic systems. *J. Heterocyclic Chem.* **1973**, *10*, 479.
2. Coburn, R. A.; Glennon, R. A. Synthesis and properties of mesoionic thiazolo[3,2-a]pyrimidin-5,7-diones. *J. Heterocyclic Chem.* **1973**, *10*, 487.
3. Coburn, R. A.; Glennon, R. A. Synthesis and *in vitro* antibacterial properties of mesoionic thiazolo[3,2-a]pyrimidin-5,7-diones and mesoionic 1,3,4-thiadiazolo[3,2a]pyrimidin-5,7-diones. *J. Pharm. Sci.* **1973**, *62*, 1785.
4. Coburn, R. A.; Bhooshan, B.; Glennon, R. A. The preparation of 2-alkylamino-1,3,4-thiadiazoles. *J. Org. Chem.* **1973**, *38*, 3947.
5. Coburn, R. A.; Glennon, R. A.; Chmielewicz, Z. *In vitro* antibacterial activity of mesoionic 1,3,4-thiadiazolo [3,2a]pyrimidin-5,7-diones. *J. Med. Chem.* **1974**, *17*, 1025.
6. Glennon, R. A. A quantum chemical investigation of the π electronic structure of the hallucinogenic N,N-dimethyltryptamines. *Res. Commun. Chem. Path. Pharmacol.* **1974**, *9*, 185.
7. Glennon, R. A.; Von Strandtmann, M. Cyclodehydration of N-heteroarylcarbamoylmethylbenzamides. *J. Heterocyclic Chem.* **1975**, *12*, 135.
8. Glennon, R. A.; Gessner, P. K. The electronic and serotonin receptor binding affinity properties of N,N-dimethyltryptamine analogs. *Res. Commun. Chem. Path. Pharmacol.* **1977**, *18*, 453.
9. Kier, L. B.; Glennon, R. A. Psychotomimetic phenalkylamines as serotonin agonists: An SAR analysis. *Life Sciences* **1978**, *20*, 1589.
10. Glennon, R. A.; Martin, B.; Johnson, K.; End, D. 7,N,N-Trimethyltryptamine: A selective inhibitor of synaptosomal serotonin uptake. *Res. Commun. Chem. Path. Pharmacol.* **1978**, *19*, 161.
11. Glennon, R. A.; Liebowitz, S. M.; Mack, E. C. Serotonin receptor binding affinities of several hallucinogenic phenalkylamine and N,N-dimethyltryptamine analogs. *J. Med. Chem.* **1978**, *21*, 822.
12. Glennon, R. A.; Kier, L. B. LSD Analogs as serotonin antagonists: A molecular connectivity SAR analysis. *Eur. J. Med. Chem.* **1978**, *13*, 219.
13. Glennon, R. A.; Rogers, M. E.; Bass, R. G.; Ryan, S. B. Mesoionic xanthine analogs as inhibitors of cyclic AMP phosphodiesterase; *J. Pharm. Sci.* **1978**, *67*, 1762.
14. Watson, H.; Glennon, R. A. Common drug interactions. *Anesthesia Update* **1978**, *6*, 1.
15. Glennon, R. A.; Kier, L. B.; Shulgin, A. T. A molecular connectivity analysis of mescaline analogs. *J. Pharm. Sci.* **1979**, *68*, 906.
16. Glennon, R. A.; Rosecrans, J. A.; Young, R.; Gaines, J. Hallucinogens as discriminative stimuli: Generalization of a 5-methoxy DMT stimulus with DOM. *Life Sciences* **1979**, *24*, 993.
17. Glennon, R. A.; Bass, R. G.; Schubert, E. Alkylation of thiazolo[3,2-a]pyrimidin-5,7-diones; *J. Heterocyclic Chem.* **1979**, *16*, 903.
18. Glennon, R. A. The effect of chirality on serotonin receptor affinity. *Life Sciences* **1979**, *24*, 1487.
19. Glennon, R. A. A similarity between rat fundus serotonin receptors and brain serotonin binding sites. *Res. Commun. Psychol. Psychiat. Behav.* **1979**, *4*, 333.
20. Glennon, R. A.; Gessner, P. K. Serotonin receptor affinities of tryptamine analogs. *J. Med. Chem.* **1979**, *22*, 428.
21. Glennon, R. A.; Gessner, P. K.; Godse, D.; Kline, B. Bufotenine esters. *J. Med. Chem.* **1979**, *22*, 1414.
22. Rosecrans, J. A.; Glennon, R. A. Drug-induced cues in studying mechanisms of drug action. *Neuropharmacology* **1979**, *18*, 981.

1980-1984

23. Glennon, R. A.; Rogers, M. E.; El-Said, M. K. Imidazo[1,2a]-pyrimidines and 1,2,4-triazolo[1,5-a]pyrimidines: Two new examples of mesoionic xanthine analogs. *J. Heterocyclic Chem.* **1980**, *17*, 337.
24. Glennon, R. A.; Liebowitz, S. M.; Anderson III, G. M. Serotonin receptor affinities of psychoactive phenalkylamine analogs. *J. Med. Chem.* **1980**, *23*, 294.

25. Minnema, D.; Krynock, G.; Young, R.; Glennon, R. A.; Rosecrans, J. A. Role of dorsal raphe neurons to the ability of LSD to act as a discriminative stimulus. *J. Alcohol Drug Abuse* **1980**, *1*, 29.
26. Glennon, R. A.; Young, R.; Rosecrans, J. A.; Kallman, M. J. Hallucinogenic agents as discriminative stimuli: A correlation with serotonin receptor affinities. *Psychopharmacology* **1980**, *68*, 155.
27. Glennon, R. A.; Liebowitz, S. M.; Doot, D. L.; Rosecrans, J. A. Demethyl analogues of psychoactive methoxy phenylisopropylamines: Synthesis and serotonin receptor affinities. *J. Med. Chem.* **1980**, *23*, 990.
28. Glennon, R. A.; Schubert, E.; Jacyno, J. M. Studies on several 7-substituted N,N-dimethyltryptamines. *J. Med. Chem.* **1980**, *23*, 1222.
29. Glennon, R. A.; Doot, D. L.; Young, R. DOM and related 2,5-dimethoxy-4-alkylphenylisopropylamines: behavioral and serotonin receptor properties. *Pharmacol. Biochem. Behav.* **1981**, *14*, 287.
30. Glennon, R. A.; Rogers, M. E.; Smith, J. D.; El-Said, M. K.; Egle, J. L. Mesoionic xanthine analogues: phosphodiesterase inhibitory and hypotensive activity. *J. Med. Chem.* **1981**, *24*, 658.
31. Glennon, R. A.; Salley, J. J.; Steinsland, O. S.; Nelson, S. Synthesis and evaluation of novel alkylpiperazines as potential dopamine antagonists. *J. Med. Chem.* **1981**, *24*, 678.
32. Glennon, R. A.; Gaines, J. J.; Rogers, M. E. Benz-fused mesoionic xanthine analogues as inhibitors of cyclic-amp phosphodiesterase. *J. Med. Chem.* **1981**, *24*, 766.
33. Rogers, M. E.; Glennon, R. A.; Smith, J. D.; Boots, M. R.; Nanavati, N.; Maconaughey, J. E.; Aub, D.; Thomas, S.; Bass, R. G.; Mbagwu, G. Mesoionic purinone analogues as inhibitors of cyclic-amp phosphodiesterase: A comparison of several ring systems. *J. Med. Chem.* **1981**, *24*, 1284.
34. Glennon, R. A.; Showalter, D. The effects of cathinone and several related derivatives on locomotor activity. *Res. Commun. Subst. Abuse* **1981**, *2*, 186.
35. Glennon, R. A.; Rosecrans, J. A. Speculations on the mechanism of action of hallucinogenic indolealkylamines. *Neurosci. Biobehav. Rev.* **1981**, *5*, 197.
36. Young, R.; Glennon, R. A.; Rosecrans, J. A. Discriminative stimulus properties of the hallucinogenic agent DOM. *Commun. Psychopharmacol.* **1981**, *4*, 501.
37. Glennon, R. A. Serotonin receptor interactions of harmaline and several related β -carbolines. *Life Sciences* **1981**, *29*, 861.
38. Glennon, R. A.; Schubert, E.; Bass, R. G. Synthesis of mesoionic xanthine nucleosides. *Tetrahedron Lett.* **1981**, *29*, 2753.
39. Domelsmith, L. N.; Eaton, T. A.; Houk, K. N.; Anderson, G. M.; Glennon, R. A.; Shulgin, A.T.; Castagnoli, N.; Kollman, P. A. Relationship between physical properties and pharmacological actions of amphetamine analogues. *J. Med. Chem.* **1981**, *24*, 1414.
40. Glennon, R. A.; Rosecrans, J. A.; Young, R. Behavioral properties of psychoactive phenylisopropylamines. *Eur. J. Pharmacol.* **1981**, *76*, 353.
41. Glennon, R. A.; Young, R.; Benington, F.; Morin, R. Hallucinogens as discriminative stimuli: A comparison of 4-OMe DMT and 5-OMe DMT with their methylthio counterparts. *Life Sciences* **1982**, *30*, 465.
42. Glennon, R. A.; Jacyno, J. M.; Salley, J. J. 2,3-Dihydro and carbocyclic analogues of tryptamines: Interaction with serotonin receptors. *J. Med. Chem.* **1982**, *25*, 68.
43. Glennon, R. A.; Rosecrans, J. A.; Young, R. Discriminative stimulus properties of DOM and several molecular modifications. *Pharmacol. Biochem. Behav.* **1982**, *16*, 553.
44. Glennon, R. A.; Young, R.; Rosecrans, J. A. A comparison of the behavioral effects of DOM homologs. *Pharmacol. Biochem. Behav.* **1982**, *16*, 557.
45. Glennon, R. A.; Liebowitz, S. M. Serotonin receptor affinity of cathinone analogs. *J. Med. Chem.* **1982**, *25*, 393.
46. Young, R.; Rosecrans, J. A.; Glennon, R. A. Comparative discriminative stimulus effects of 5-methoxy-n,n-dimethyltryptamine and LSD. *Life Sciences* **1982**, *30*, 2057.
47. Glennon, R. A.; Young, R.; Rosecrans, J. A.; Anderson, G. Discriminative stimulus properties of MDA analogs. *Biol. Psychiat.* **1982**, *17*, 807.

48. Salley, J. J.; Glennon, R. A. Studies on simplified ergoline derivatives. A general six-step synthesis of phenyl-substituted 4-methyl-3,4a,5,6,10b-hexahydrobenzo[f]quinolin-1(2H)-one analogs. *J. Heterocyclic Chem.* **1982**, *19*, 545.
49. Glennon, R. A.; Young, R. A comparison of the behavioral properties of di- and trimethoxyphenylisopropylamines. *Pharmacol. Biochem. Behav.* **1982**, *17*, 603.
50. Glennon, R. A.; Young, R.; Benington, F.; Morin, R. D. Behavioral and serotonin receptor properties of 4-substituted derivatives of the hallucinogen 1-(2,5-dimethoxyphenyl)-2-aminopropane. *J. Med. Chem.* **1982**, *25*, 1163.
51. Glennon, R. A.; Rosecrans, J. Indolealkylamine and phenalkylamine hallucinogens: A brief overview. *Neurosci. Biobehav. Rev.* **1982**, *6*, 489.
52. Lund, M. Q.; Kier, L. B.; Glennon, R. A.; Egle, J. L. Preliminary studies of mesoionic 3-(substituted-aryl)- ψ -oxatriazoles as potential antihypertensive agents. *J. Med. Chem.* **1982**, *25*, 1503.
53. Kline, T. B.; Benington, F.; Morin, R. D.; Beaton, J. M.; Glennon, R. A.; Domelsmith, L.; Houk, K. N.; Rozeboom, M. D. Structure-activity relationships for hallucinogenic N,N-dialkyltryptamines: Photoelectron spectra and 5-HT receptor affinities of methylthio and methylenedioxy derivatives. *J. Med. Chem.* **1982**, *25*, 1381.
54. Glennon, R. A.; Young, R.; Jacyno, J. M. Indolealkylamine and phenalkylamine hallucinogens: Effect of α -methyl and n-methyl substituents on behavioral activity. *Biochem. Pharmacol.* **1983**, *32*, 1267.
55. Glennon, R. A.; Young, R.; Jacyno, J. M.; Slusher, R. M.; Rosecrans, J. A. DOM-stimulus generalization to LSD and other hallucinogenic indolealkylamines. *Eur. J. Pharmacol.* **1983**, *86*, 453.
56. Glennon, R. A.; Jacyno, J. M.; Young, R. A comparison of the behavioral properties of (\pm), (+) and (-)-5-methoxy- α -methyltryptamine. *Biol. Psychiat.* **1983**, *18*, 493.
57. Young, R.; Rosecrans, J. A.; Glennon, R. A. Behavioral effects of 5-methoxy-N,N-dimethyltryptamine and dose-dependent antagonism by BC-105. *Psychopharmacology* **1983**, *80*, 156.
58. Schubert, E. M.; Bass, R. G.; Glennon, R. A. Synthesis of mesoionic xanthine nucleosides. *Nucleosides/Nucleotides* **1983**, *2*, 127.
59. Glennon, R. A.; Rosecrans, J. A.; Young, R. Drug-induced discrimination: A description of the paradigm and a review of its specific application to the study of hallucinogenic agents. *Med. Res. Rev.* **1983**, *3*, 289.
60. Glennon, R. A.; Young, R.; Rosecrans, J. A. Antagonism of the effects of the hallucinogen DOM and the purported serotonin agonist quipazine by 5-HT₂ antagonists. *Eur. J. Pharmacol.* **1983**, *91*, 189.
61. Glennon, R. A.; Young, R.; Jacyno, J. M.; Nelson, D. Synthesis and evaluation of a novel series of N,N-di methylisotryptamines. *J. Med. Chem.* **1984**, *27*, 41.
62. Mbagwu, G.; Bass, R. G.; Glennon, R. A. Carbon-13 nuclear magnetic resonance spectra of some mesoionic xanthine analogs. *Org. Magnetic Res.* **1983**, *21*, 527.
63. Glennon, R. A.; Young, R. MDA: A psychoactive agent with dual stimulus effects. *Life Sciences* **1984**, *34*, 379.
64. Glennon, R. A.; Young, R. Further investigation of the discriminative stimulus properties of MDA. *Pharmacol. Biochem. Behav.* **1984**, *20*, 501.
65. Wood, P. L.; Pilapil, C.; LaFaille, F.; Nair, N. P.; Glennon, R. A. Unique [³H]tryptamine binding sites in rat brain: Distribution and pharmacology. *Arch. Int. Pharmacodyn.* **1984**, *268*, 194.
66. Glennon, R. A.; Young, R.; Soine, W. 1-(2,3-Methylenedioxyphenyl)-2-aminopropane: A preliminary investigation. *Gen. Pharmacol.* **1984**, *15*, 361.
67. Aceto, M.; Rosecrans, J. A.; Young, R.; Glennon, R. A. Similarity between (+)-amphetamine and amfonelic acid. *Pharmacol. Biochem. Behav.* **1984**, *20*, 635.
68. Young, R.; Dewey, W. L.; Glennon, R. A. Stereoselective stimulus effects of 3-methylflunitrazepam and pentobarbital. *Life Sciences* **1984**, *34*, 1977-1983.
69. Schechter, M. D.; Rosecrans, J. A.; Glennon, R. A. Comparison of the behavioral effects of cathinone, amphetamine and apomorphine. *Pharmacol. Biochem. Behav.* **1984**, *20*, 181.

70. Glennon, R. A.; Schechter, M. D.; Rosecrans, J. A. Discriminative stimulus properties of *S*(-) and *R*(+)-cathinone, (+)-cathine, and several structural modifications. *Pharmacol. Biochem. Behav.* **1984**, - 21, 1.
71. Glennon, R. A.; Young, R. MDA: An agent that produces stimulus effects similar to those of 3,4-DMA, LSD and Cocaine. *Eur. J. Pharmacol.* **1984**, 99, 249.
72. Shannon, M.; Battaglia, G.; Glennon, R. A.; Titeler, M. 5-HT₁ and 5-HT₂ receptor binding properties of derivatives of the hallucinogen 1-(2,5-dimethoxyphenyl)-2-aminopropane. *Eur. J. Pharmacol.* **1984**, 102, 23.
73. Glennon, R. A.; Hauck, A.; McKenney, J. D.; Young, R. Structure-activity relationships of amphetamine analogs using drug discrimination methodology. *Pharmacol. Biochem. Behav.* **1984**, 21, 895.
74. Glennon, R. A.; Tejani, S. Mesoionic nucleosides: Studies on mesoionic imidazo[2,1-b]-1,3-thiazine derivatives. *Nucleosides/Nucleotides* **1984**, 3, 389.
75. Glennon, R. A.; Tejani, S.; Padgett, W.; Daly, J. W. Mesoionic xanthine analogues as antagonists of adenosine receptors. *J. Med. Chem.* **1984**, 27, 1364.
76. Glennon, R. A.; Titeler, M.; McKenney, J. D. Evidence for 5-HT₂ involvement in the mechanism of action of hallucinogenic agents. *Life Sci.* **1984**, 35, 2505.
77. Glennon, R. A.; McKenney, J. D.; Young, R. Discriminative stimulus properties of the serotonin agonist 1-(3-trifluoromethylphenyl)piperazine (TFMPP). *Life Sci.* **1984**, 35, 1475.

1985-1989

78. Glennon, R. A.; Young, R.; Hauck, A. E., Structure-activity studies on methoxy-substituted phenylisopropylamines using drug discrimination methodology. *Pharmacol. Biochem. Behav.* **1985**, 22, 723.
79. Titeler, M.; Herrick, C.; Lyon, R. A.; McKenney, J. D.; Glennon, R. A. [³H]DOB: A specific radioligand for 5-HT₂ receptors. *Eur. J. Pharmacol.* **1985**, 117, 145.
80. Young, R.; Glennon, R. A.; Brase, D.; Dewey, W. L. Potencies of diazepam metabolites in generalizing to the diazepam discriminative stimulus. *Life Sci.* **1985**, 39, 17.
81. Glennon, R. A.; Hauck, A. E. Mechanistic studies on DOM as a discriminative stimulus. *Pharmacol. Biochem. Behav.* **1985**, 23, 937.
82. Schubert, E. M.; Schram, K. H.; Glennon, R. A. Mass spectrometry of modified nucleic acid bases and nucleosides. Class II mesoionic nucleosides and bases derived from thiazolo[3,2-a]pyrimidine-5,7-diones. *J. Heterocyclic Chem.* **1985**, 22, 889.
83. Schechter, M. D.; Glennon, R. A. Cathinone, cocaine and methamphetamine: Similarity of behavioral effects. *Pharmacol. Biochem. Behav.* **1985**, 22, 913.
84. Mbagwu, G. O.; Bass, R. G.; Glennon, R. A. Studies of amine-induced ring-opening of some mesoionic xanthenes. *J. Heterocyclic Chem.* **1985**, 22, 465.
85. Glennon, R. A.; McKenney, J. D.; Lyon, R. A. Titeler, M. 5-HT₁ and 5-HT₂ binding characteristic of 1-(2,5-dimethoxy-4-bromophenyl)-2-aminopropane analogs. *J. Med. Chem.* **1986**, 29, 194.
86. McKenney, J. D.; Glennon, R. A. TFMPP may produce its stimulus effects via a 5-HT_{1B} mechanism. *Pharmacol. Biochem. Behav.* **1986**, 24, 43.
87. Lyon, R. A.; Titeler, M.; McKenney, J. D.; Magee, P. S.; Glennon, R. A. Synthesis and evaluation of phenyl- and benzoylpiperazines as potential serotonergic agents. *J. Med. Chem.* **1986**, 29, 630.
88. Young, R.; Glennon, R. A. Amphetamine and related phenalkylamines as discriminative stimuli: A review of the literature. *Med. Res. Rev.* **1986**, 6, 99.
89. Lyon, R. A.; Glennon, R. A.; Titeler, M. 3,4-Methylenedioxymethamphetamine (MDMA): Stereoselective interactions at brain 5-HT₁ and 5-HT₂ sites. *Psychopharmacology* **1986**, 88, 525.
90. Glennon, R. A. Discriminative stimulus properties of the 5-HT_{1A} agonist 8-OH DPAT. *Life Sci.* **1986**, 39, 825.
91. Glennon, R. A. Discriminative stimulus properties of the serotonergic agent 1-(2,5-dimethoxy-4-iodophenyl)-2-aminopropane (DOI). *Pharmacol. Biochem. Behav.* **1986**, 25, 135.

92. Kalix, P.; Glennon, R. A. Further evidence for an amphetamine-like mechanism of action of the alkaloid cathinone. *Biochem. Pharmacol.* **1986**, *35*, 3015.
93. Glennon, R. A. Discriminative stimulus properties of phenylisopropylamine derivatives. *Drug Alcohol Depend.* **1986**, *17*, 119.
94. Glennon, R. A.; Titeler, M.; Young, R. Structure-activity relationships and mechanism of action of hallucinogenic agents based on drug discrimination and radioligand binding studies. *Psychopharmacol. Bull.* **1986**, *22*, 953-958.
95. Glennon, R. A.; Slusher, R. M.; Lyon, R. A.; Titeler, M.; McKenney, J. D. 5-HT₁ and 5-HT₂ binding characteristics of some quipazine analogues. *J. Med. Chem.* **1986**, *29*, 2375-2380.
96. Young, R.; Rosecrans, J. A.; Glennon, R. A. Further studies on the dose-dependent stimulus properties of 5-methoxy-N,N-dimethyltryptamine. *Pharmacol. Biochem. Behav.* **1986**, *25*, 1207-1210.
97. Rasmussen, K.; Glennon, R. A.; Aghajanian, G. K. Phenethylamine hallucinogens in the locus coeruleus: potency of action correlates with rank order of 5-HT₂ binding affinity. *Eur. J. Pharmacol.* **1986**, *132*, 79-82.
98. Glennon, R. A. Central serotonin receptors as targets for drug research. *J. Med. Chem.* **1987**, *30*, 1-12.
99. Glennon, R. A.; Little, P. J.; Rosecrans, J. R.; Yousif, M. Y. The effect of MDMA ("Ecstasy") and its optical isomers on schedule-controlled responding in mice. *Pharmacol. Biochem. Behav.* **1987**, *26*, 425.
100. Glennon, R. A.; Titeler, M.; Seggel, M. R.; Lyon, R. A. N-Methyl derivatives of the 5-HT₂ agonist 1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane. *J. Med. Chem.* **1987**, *30*, 930-932.
101. Glennon, R. A.; Yousif, M. Y.; Naiman, N.; Kalix, P. Methcathinone: A new and potent amphetamine-like agent. *Pharmacol. Biochem. Behav.* **1987**, *26*, 547-551.
102. Kerdawy, M.; Bayomi, S.; Shehata, I.; Glennon, R. A. Condensed 1,2,3-triazines: 1,3,4-thiadiazolo[3,2-a]-1,3,5-triazines and isoxazolo[2,3-a]-1,3,5-triazines. *J. Heterocyclic Chem.* **1987**, *24*, 501-504.
103. Rosecrans, J. A.; Glennon, R. A. The effect of MDA and MDMA ("Ecstasy") isomers in combination with piperone on operant responding in mice. *Pharmacol. Biochem. Behav.* **1987**, *28*, 39-42.
104. Shehata, I.; Glennon, R. A. Mesoionic isoxazolo[2,3-a]pyrimidindiones and 1,3,4-oxadiazolo[3,2-a]pyrimidindiones as potential adenosine antagonists. *J. Heterocyclic Chem.* **1987**, *24*, 1291-1295.
105. Seggel, M. R.; Qureshi, G. D.; Glennon, R. A. Effect of 5-HT₂-selective agonists on cat platelet aggregation. *Life Sci.* **1987**, *41*, 1077-1081.
106. Titeler, M.; Lyon, R. A.; Davis, K. H.; Glennon, R. A. Selectivity of serotonergic drugs for multiple brain 5-HT receptors: Role of [³H]DOB, a 5-HT₂ agonist radioligand. *Biochem. Pharmacol.* **1987**, *36*, 3265-3271.
107. Young, R.; Glennon, R. A.; Dewey, W. L.; Effects of pyrazolopyridines and a triazolopyridazine on a pentobarbital discriminative stimulus. *Psychopharmacology* **1987**, *93*, 494-497.
108. Young, R.; Glennon, R. A. Stimulus properties of benzodiazepines: Correlation with binding affinities, therapeutic potencies, and structure-activity relationships. *Psychopharmacology* **1987**, *93*, 529-533.
109. Glennon, R. A.; Seggel, M. R.; Soine, W. H.; Lyon, R. A.; Davis, K.; Titeler, M. [¹²⁵I]-1-(2,5-Dimethoxy-4-iodophenyl)-2-aminopropane: An iodinated radioligand that specifically labels the agonist high-affinity state of 5-HT₂ serotonin receptors. *J. Med. Chem.* **1988**, *31*, 5-7.
110. Titeler, M.; Lyon, R. A.; Glennon, R. A. Radioligand binding evidence implicates the brain 5-HT₂ receptor as a site of action for LSD and phenylisopropylamine hallucinogens. *Psychopharmacology* **1988**, *94*, 213-216.
111. Lyon, R. A.; Titeler, M.; Seggel, M. R.; Glennon, R. A. Indolealkylamine analogs share 5-HT₂ binding characteristics with phenalkylamine hallucinogens. *Eur. J. Pharmacol.* **1988**, *145*, 291-296.
112. Glennon, R. A.; Pierson, M. E.; McKenney, J. D. Stimulus generalization of 1-(3-trifluoromethylphenyl)piperazine (TFMPP) to propranolol, pindolol, and mesulergine. *Pharm. Biochem. Behav.* **1988**, *29*, 197-199.
113. Glennon, R. A.; Yousif, M.; Patrick, G. Stimulus properties of 1-(3,4-methylenedioxyphenyl)-2-aminopropane (MDA) analogs. *Pharmacol. Biochem. Behav.* **1988**, *29*, 443-449.

114. Glennon, R. A.; Titeler, M.; Lyon, R. A.; Slusher, R. M. N,N-Di-*n*-propylserotonin: Binding at serotonin sites and a comparison with 8-hydroxy-2-(di-*n*-propylamino)tetratin. *J. Med. Chem.* **1988**, *31*, 867-869.
 115. Kalix, P.; Yousif, M. Y.; Glennon, R. A. Differential effect of the enantiomers of methylenedioxymphetamine (MDMA) on the release of radioactivity from [³H]dopamine-prelabeled rat striatum. *Res. Commun. Substance Abuse* **1988**, *9*, 45-51.
 116. Glennon, R. A.; Titeler, M.; Lyon, R. A. A preliminary investigation of the psychoactive agent 4-bromo-2,5-dimethoxyphenethylamine: A potential drug of abuse. *Pharmacol. Biochem. Behav.* **1988**, *30*, 597-601.
 117. Glennon, R. A.; Naiman, N. A.; Lyon, R. A.; Titeler, M. Arylpiperazine derivatives as high-affinity serotonin 5-HT_{1A} ligands. *J. Med. Chem.* **1988**, *31*, 1968-1971.
 118. Glennon, R. A.; Naiman, N. A.; Pierson, M. E.; Titeler, M.; Lyon, R. A. NAN-190: An arylpiperazine analog that antagonizes the stimulus effects of the 5-HT_{1A} agonist 8-hydroxy-2-(di-*n*-propylamino)tetratin. *Eur. J. Pharmacol.* **1988**, *154*, 339-341.
 119. Glennon, R. A.; Ismaiel, A. M.; Martin, B.; Poff, D.; Sutton, M. A preliminary behavioral investigation of PMMA, the 4-methoxy analog of methamphetamine. *Pharmacol. Biochem. Behav.* **1988**, *31*, 9-13.
 120. Glennon, R. A.; Naiman, N. A.; Pierson, M. E.; Titeler, M.; Lyon, R. A.; Herndon, J. L.; Misenheimer, B. Stimulus properties of arylpiperazines: NAN-190, a potential 5-HT_{1A} serotonin antagonist. *Drug Dev. Res.* **1989**, *16*, 335-343.
 121. Naiman, N. A.; Lyon, R. A.; Bullock, A. E.; Rydeleck, L. T.; Titeler, M.; Glennon, R. A. 2-(Alkylamino)tetratin derivatives: Interaction with 5-HT_{1A} serotonin binding sites. *J. Med. Chem.* **1989**, *32*, 253-256.
 122. Pierson, M. E.; Lyon, R. A.; Titeler, M.; Kowalski, P.; Glennon, R. A. Design and synthesis of propranolol analogues as serotonergic agents. *J. Med. Chem.* **1989**, *32*, 859-863.
 123. Fitzgerald, R. L.; Blanke, R. V.; Rosecrans, J. A.; Glennon, R. A. Stereochemistry of the metabolism of MDMA to MDA. *Life Sci.* **1989**, *45*, 295-301.
 124. Sadzot, B.; Baraban, J. M.; Glennon, R. A.; Lyon, R. A.; Leonhardt, S.; Jan, C-R.; Titeler, M. Hallucinogenic drug interactions at human brain 5-HT₂ receptors: Implications for treating LSD induced hallucinogenesis. *Psychopharmacology* **1989**, *98*, 495-499.
 125. Glennon, R. A.; Naiman, N. A.; Pierson, M. E.; Smith, J. D.; Ismaiel, A. M.; Titeler, M.; Lyon, R. A. N-(Phthalimidoalkyl) derivatives of serotonergic agents: A common interaction at 5-HT_{1A} serotonin binding sites? *J. Med. Chem.* **1989**, *32*, 1921-1926.
 126. Glennon, R. A.; Misenheimer, B. Stimulus effects of N-mono ethyl-(3,4-methylenedioxyphenyl)-2-aminopropane (MDE) and N-hydroxy-1-(3,4-methylenedioxyphenyl)-2-aminopropane (N-OH MDA) in rats trained to discriminate MDMA from saline. *Pharmacol. Biochem. Behav.* **1989**, *33*, 900-912.
 127. Glennon, R. A.; Ismaiel, A. M.; McCarthy, B. G.; Peroutka, S. J. Binding of arylpiperazines to 5-HT₃ serotonin receptors: Results of a structure-affinity study. *Eur. J. Pharmacol.* **1989**, *168*, 387-392.
 128. Ismaiel, A. M.; Titeler, M.; Glennon, R. A. γ -Carboline derivatives as potential serotonergic agents. *Mansoura J. Pharm. (Egypt)* **1989**, *6*, 1-13.
 129. Fitzgerald, R. L.; Blanke, R. V.; Glennon, R. A.; Yousif, M.; Rosecrans, J. A.; Poklis, A. Determination of 3,4-methylenedioxymphetamine (MDA) and N-methyl-3,4-methylenedioxymphetamine (MDMA) enantiomers in whole blood. *J. Chromatog.* **1989**, *490*, 59-69.
- 1990-1994**
130. Ismaiel, A. M.; Titeler, M.; Miller, K. J.; Smith, T. S.; Glennon, R. A. 5-HT₁ and 5-HT₂ binding profiles of the serotonergic agents α -methylserotonin and 2-methylserotonin. *J. Med. Chem.* **1990**, *33*, 755-758.
 131. Glennon, R. A. Serotonin receptors: Clinical Implications. *Neurosci. Biobehav. Rev.* **1990**, *14*, 35-47.
 132. Kerdawy, M. M.; Ismaiel, A. M.; Gineinah, M.; Glennon, R. A. A convenient synthesis of 3-aryl-1,2,4-triazolo [4,3-*c*]quinazolines. *J. Heterocyclic Chem.* **1990**, *27*, 497-501.

133. Seggel, M. R.; Yousif, M. Y.; Lyon, R. A.; Titeler, M.; Roth, B. L.; Suba, E. A.; Glennon, R. A. An SAR study of the binding of 4-substituted analogues of 1-(2,5-dimethoxyphenyl)-2-aminopropane at 5-HT₂ serotonin receptors. *J. Med. Chem.* **1990**, *33*, 1032-1036.
134. Gineinah, M.; Ismaiel, A. M.; Kerdawy, M.; Glennon, R. A. Mesoionic 1,2,4-triazolo[4,3-c]quinazolines. *J. Heterocyclic Chem.* **1990**, *27*, 723-726.
135. Glennon, R. A.; Misenheimer, B. Stimulus properties of a new designer drug: 4-methylaminorex ("U4Euh"). *Pharmacol. Biochem. Behav.* **1990**, *35*, 517-521.
136. Darmani, N.; Martin, B. R.; Pandey, U.; Glennon, R. A. Do functional relationships exist between 5-HT_{1A} and 5-HT₂ receptors? *Pharmacol. Biochem. Behav.* **1990**, *36*, 901-906.
137. Higgs, R.; Glennon, R. A. Stimulus properties of ring-methyl amphetamine analogs. *Pharmacol. Biochem. Behav.* **1990**, *37*, 835-837.
138. Glennon, R. A.; Chaurasia, C.; Titeler, M. Binding of indolylalkylamines at 5-HT₂ sites: Examination of a hydrophobic binding region. *J. Med. Chem.* **1990**, *33*, 2777-2784.
139. Glennon, R. A.; Raghupathi, R. K. Serotonin receptor ligands. *Current CNS Patents* **1990**, *1*, 323-343.
140. Glennon, R. A.; Battaglia, G.; Smith, J. (-)PPAP: A new selective ligand for σ binding sites. *Pharmacol. Biochem. Behav.* **1990**, *37*, 557-559.
141. Darmani, N. A.; Martin, B. R.; Glennon, R. A. Withdrawal from chronic treatment with (\pm)DOI causes both sub- and supersensitivity to 5-HT₂ receptor-induced head-twitch behavior in mice. *Eur. J. Pharmacol.* **1990**, *186*, 115-118.
142. Darmani, N. A.; Martin, B. R.; Pandey, U.; Glennon, R. A. Pharmacological classification of ear-scratch response in mice as a behavioral model for selective 5-HT₂ receptor agonists and evidence for 5-HT_{1B}- and 5-HT₂-receptor interactions. *Pharmacol. Biochem. Behav.* **1990**, *37*, 95-99.
143. Rydelek-Fitzgerald, L.; Teitler, M.; Ismaiel, A. M.; Glennon, R. A. NAN-190: Agonist and antagonist interactions with brain 5-HT_{1A} receptors. *Brain Res.* **1990**, *532*, 191-196.
144. Yousif, M. Y.; Fitzgerald, R. L.; Narasimhachari, N.; Rosecrans, J. A.; Glennon, R. A. Identification of metabolites of 3,4-methylenedioxymethamphetamine in rats. *Drug Alcohol Depend.* **1990**, *26*, 127-135.
145. Appel, N. M.; Mitchell, W. M.; Garlick, R. K.; Glennon, R. A.; Teitler, M.; De Souza, E. B. Autoradiographic characterization of (\pm)-1-(2,5-dimethoxy-4-[¹²⁵I]iodophenyl)-2-aminopropane ([¹²⁵I]DOI) binding to 5-HT₂ and 5-HT_{1C} receptors in rat brain. *J. Pharmacol. Exp. Ther.* **1990**, *255*, 843-857.
146. Glennon, R. A. Do hallucinogens act as 5-HT₂ agonists? *Neuropsychopharmacology* **1990**, *3*, 509-517.
147. Glennon, R. A.; DeVry, J.; Spencer, D. G.; Glaser, T. Stimulus properties of tifenaridine: A novel antidepressant agent. *Pharmacol. Biochem. Behav.* **1990**, *37*, 769-771.
148. Glennon, R. A.; Smith, J. D.; Ismaiel, A. M.; El-Ashmawy, M.; Battaglia, G.; Fischer, J. B. Identification and exploitation of the sigma opiate pharmacophore. *J. Med. Chem.* **1991**, *34*, 1094-1098.
149. Darmani, N.; Martin, B. B.; Pandey, U.; Glennon, R. A. Inhibition of 5-HT₂-mediated head-twitch response by cocaine via indirect stimulation of adrenergic α_2 and serotonergic 5-HT_{1A} receptors. *Pharmacol. Biochem. Behav.* **1991**, *38*, 353-357.
150. Glennon, R. A. Serotonin receptors and site-selective agents. *J. Physiol. Pharmacol. (Poland)* **1991**, *1*, 42-60.
151. Glennon, R. A.; Ismaiel, A. M.; Chaurasia, C.; Titeler, M. 5-HT_{1D} serotonin receptors: Results of a structure-affinity investigation. *Drug Dev. Res.* **1991**, *22*, 25-36.
152. Glennon, R. A.; Ismaiel, A.; Smith, J. D.; Yousif, M. Y.; El-Ashmawy, M.; Herndon, J. L.; Fischer, J. B.; Burke Howie, K. J.; Server, A. A. Binding of substituted and conformationally restricted derivatives of N-(3-phenyl-*n*-propyl)-1-phenyl-2-aminopropane at σ receptors. *J. Med. Chem.* **1991**, *34*, 1855-1859.
153. Glennon, R. A.; El-Ashmawy, M.; Ismaiel, A. M.; Fischer, J. B. Binding of N-substituted 2-phenylaminoethanes to σ and serotonin 5-HT_{1A} receptors. *Med. Chem. Res.* **1991**, *1*, 109-114.
154. Raghupathi, R. K.; Rydelek-Fitzgerald, L.; Teitler, M.; Glennon, R. A. Analogues of the 5-HT_{1A} serotonin antagonist 1-(2-methoxyphenyl)-4-[4-(2-phthalimido)-butyl]piperazine with reduced α_1 -adrenergic affinity. *J. Med. Chem.* **1991**, *34*, 2633-2638.

155. Glennon, R. A.; Bartyzel, P.; Teitler, M. Binding of benz[e]- and benz[g]-fused tryptamine derivatives at serotonin receptors: Evidence for a region of bulk tolerance. *Med. Chem. Res.* **1991**, *1*, 201-206.
156. Glennon, R. A.; El-Ashmawy, M.; Fischer, J. B.; Burke Howie, K. J.; Ismaiel, A. M. N-Substituted 5-phenyl pentylamines: A new class of σ receptor ligands. *Med. Chem. Res.* **1991**, *1*, 207-212.
157. Glennon, R. A.; Darmani, N.; Martin, W. Modulation of the behavioral effects of serotonergic agents by multiple 5-HT receptors *Life Sci.* **1991**, *48*, 2493-2498.
158. Glennon, R. A.; Dukat, M. Serotonin Receptors and their Ligands: A lack of selective agents. *Pharmacol. Biochem. Behav.* **1991**, *40*, 1009-1017.
159. Glennon, R. A.; Yousif, M. Y.; Ismaiel, A. M.; El-Ashmawy, M. B.; Herndon, J. L.; Fischer, J. B.; Server, A. C.; Burke-Howie, K. J. Novel 1-phenylpiperazines and 4-phenylpiperidines as high-affinity sigma ligands. *J. Med. Chem.* **1991**, *34*, 3360-3365.
160. Dukat, M.; Miller, K.; Teitler, M.; Glennon, R. A. Binding of amine-substituted and quaternary amine analogs of serotonin at 5-HT₂ serotonin receptors. *Med. Chem. Res.* **1991**, *1*, 271-276.
161. Bakthavachalam, V.; Fell, B.; Teitler, M.; Glennon, R. A.; Neumeyer, J. L. Fluorescent probes for serotonin 5-HT_{1A} receptors: synthesis, receptor affinity and selectivity. *Med. Chem. Res.* **1991**, *1*, 265-270.
162. Westkaemper, R. B.; Glennon, R. A. Approaches to molecular modeling studies and specific application to serotonin ligands and receptors. *Pharmacol. Biochem. Behav.* **1991**, *40*, 1019-1031.
163. Westkaemper, R. B.; Dukat, M.; Glennon, R. A. Molecular modeling of drug-receptor interactions using a 5-HT₂ receptor model. *Med. Chem. Res.* **1991**, *1*, 401-408.
164. Ablordeppey, S. Y.; El-Ashmawy, M. B.; Glennon, R. A. Analysis of the structure-activity relationships of sigma receptor ligands. *Med. Chem. Res.* **1991**, *1*, 425-438.
165. Glennon, R. A. Preface for special serotonin issue of *Pharmacol. Biochem. Behav.* **1991**, *40*, 1007.
166. Glennon, R. A.; Raghupathi, R.; Bartyzel, P.; Teitler, M.; Leonhardt, S. Binding of phenylalkylamine derivatives at 5-HT_{1C} and 5-HT₂ serotonin receptors: Evidence for a Lack of selectivity. *J. Med. Chem.* **1992**, *35*, 734-740.
167. Glennon, R. A.; Young, R.; Dukat, M. The 5-HT₂ agonist 2-methylserotonin as a training drug in drug discrimination studies. *Pharmacol. Biochem. Behav.* **1992**, *41*, 361-364.
168. El-Bermawy, M.; Raghupathi, R.; Ingher, S. P.; Teitler, M.; Maayani, S.; Glennon, R. A. 4-[4-(1-Noradamantane carboxamido)butyl]-1-(2-methoxyphenyl) piperazine: A high affinity 5-HT_{1A}-selective agent. *Med. Chem. Res.* **1992**, *2*, 88-95.
169. El-Ashmawy, M.; Ablordeppey, S. Y.; Issa, H.; Gad, L.; Fischer, J. B.; Burke Howie, K. J.; Glennon, R. A. Further investigation of 5-phenylpentylamine derivatives as novel σ receptor ligands. *Med. Chem. Res.* **1992**, *2*, 119-126.
170. El-Bermawy, M.; Lotter, H.; Glennon, R. A. Comparative molecular field analysis of the binding of arylpiperazines at 5-HT_{1A} serotonin receptors. *Med. Chem. Res.* **1992**, *2*, 290-297.
171. Ablordeppey, S. Y.; El-Ashmawy, M. B.; Glennon, R. A. Analysis of the structure-activity relationships of sigma receptor ligands. *Med. Chem. Res.* **1992**, *2*, 425-438.
172. Herndon, J. L.; Pierson, M. E.; Glennon, R. A. Mechanistic investigation of the stimulus properties of 1-(3-trifluoromethylphenyl)piperazine. *Pharmacol. Biochem. Behav.* **1992**, *43*, 739-748.
173. Glennon, R. A.; Higgs, R. Investigation of MDMA-related agents in rats trained to discriminate MDMA from saline. *Pharmacol. Biochem. Behav.* **1992**, *43*, 759-763.
174. Glennon, R. A. Concepts for design of 5-HT_{1A} serotonin agonists and antagonists. *Drug Dev. Res.* **1992**, *26*, 251-274.
175. Ablordeppey, S. Y.; Fischer, J. B.; Burke Howie, K. J.; Glennon, R. A. Design, synthesis and binding of sigma receptor ligands derived from butaclamol. *Med. Chem. Res.* **1992**, *2*, 368-375.
176. Herndon, J. L.; Ismaiel, A.; Ingher, S.; Teitler, M.; Glennon, R. A. Ketanserin analogues: Structure-affinity relationships for 5-HT₂ and 5-HT_{1C} serotonin receptor binding. *J. Med. Chem.* **1992**, *35*, 4903-4910.

177. Glennon, R. A.; Higgs, R.; Young, R.; Issa, H. Further studies on N-methyl-1-(3,4-methylenedioxypheyl)-2-aminopropane as a discriminative stimulus: Antagonism by 5-hydroxytryptamine antagonists. *Pharmacol. Biochem. Behav.* **1992**, *43*, 1099-1106.
 178. Westkaemper, R. B.; Yousif, M.; Teitler, M.; Glennon, R. A. Binding of aporphines at 5-HT_{1A} serotonin receptors: A computational investigation of potential binding modes. *Med. Chem. Res.* **1992**, *2*, 482-490.
 179. Glennon, R. A.; Dukat, M. 5-HT receptor ligands: Update 1992. *Current Drugs* **1992**, *1*, 1-45.
 180. Young, R.; Glennon, R. A. Zacopride and its optical isomers produce stereoselective antagonism of a 2-methyl serotonin discriminative stimulus. *Eur. J. Pharmacol.* **1992**, *212*, 117-119.
 181. Glennon, R. A.; Maarouf, R.; Fahmy, S.; Martin, B.; Fan, F.; Yousif, M.; Shafik, R.; Dukat, M. Structure-affinity relationships of simple nicotine analogs. *Med. Chem. Res.* **1993**, *2*, 546-551.
 182. Young, R.; Glennon, R. A. Cocaine-stimulus generalization to two new designer drugs: Methcathinone and 4-methylaminorex. *Pharmacol. Biochem. Behav.* **1993**, *45*, 229-231.
 183. Glennon, R. A.; Young, R.; Herndon, J. L. Antagonism of a (+)N-allylnormetazocine stimulus by (-)-PPAP and several structurally related analogs. *Pharmacol. Biochem. Behav.* **1993**, *45*, 865-869.
 184. Ablordeppey, S. Y.; Issa, H.; Fischer, J. B.; Burke Howie, K. J.; Glennon, R. A. Synthesis and structure-affinity relationship studies of sigma ligands related to haloperidol. *Med. Chem. Res.* **1993**, *3*, 131-138.
 185. Ismaiel, A. M.; De Los Angeles, J.; Teitler, M.; Ingher, S.; Glennon, R. A. Antagonism of 1-(2,5-dimethoxy-4-methylphenyl)-2-aminopropane stimulus with a newly identified 5-HT₂- versus 5-HT_{1C}-selective antagonist. *J. Med. Chem.* **1993**, *36*, 2519-2525.
 186. Glennon, R. A. MDMA-like stimulus effects of α -ethyltryptamine and the α -ethyl homolog of DOM. *Pharmacol. Biochem. Behav.* **1993**, *46*, 459-462.
 187. Westkaemper, R. B. and Glennon, R. A. Molecular graphics models of members of the 5-HT₂ subfamily: 5-HT_{2A}, 5-HT_{2B}, and 5-HT_{2C} receptors. *Med. Chem. Res.* **1993**, *3*, 317-334.
 188. Abou-Gharbia, M.; Ablordeppey, S. Y.; Glennon, R. A. Sigma receptors and their ligands: The Sigma Enigma. *Ann. Rep. Med. Chem.* **1993**, *28*, 1-10.
 189. Glennon, R. A.; Westkaemper, R. B. 5-HT_{1D} Receptors: A serotonin receptor population for the 1990s. *Drug News Perspect.* **1993**, *6*, 390-405.
 190. Dukat, M.; Damaj, M. I.; Glassco, W.; Dumas, D.; May, E. L.; Martin, B. R.; Glennon, R. A. Epibatidine: A very high affinity nicotine-receptor ligand. *Med. Chem. Res.* **1994**, *4*, 131-139.
 191. Damaj, M. I.; Glennon, R. A.; Martin, B. R. Involvement of the serotonergic system in the hypoactive and antinociceptive effects of nicotine in mice. *Brain Res.* **1994**, *33*, 199-203.
 192. Glennon, R. A.; Ablordeppey, S. Y.; Ismaiel, A. M.; El-Ashmawy, M. B.; Fischer, J. B.; Burke-Howie, K. Structural features important sigma-1 (σ_1) receptor binding. *J. Med. Chem.* **1994**, *37*, 1214-1219.
 193. Carroll, F. I.; Glennon, R. A.; Johnson, M. R.; Teitler, M.; Zimmerman, D. M. Hallucinogenic Agents: Drugs of abuse as neurochemical tools. *NIDA Res. Mono.* **1994**, *140*, 94-98.
 194. Glennon, R. A.; Dukat, M.; El-Bermawy, M.; Law, H.; De Los Angeles, J.; Teitler, M.; King, A.; Herrick-Davis, K. Influence of amine substituents on 5-HT_{2A} versus 5-HT_{2C} binding of phenylalkyl- and indolylalkylamines. *J. Med. Chem.* **1994**, *37*, 1929-1935.
 195. Dukat, M.; Herrick-Davis, K.; Teitler, M.; Glennon, R. A. Piperidine derivatives of serotonin as selective 5-HT_{1A} serotonin receptor agonists. *Med. Chem. Res.* **1994**, *4*, 254-258.
 196. Glennon, R. A.; Hong, S.-S.; Dukat, M.; Teitler, M. and Davis, K. 5-(Nonyloxy)tryptamine: A novel high-affinity 5-HT_{1D} serotonin receptor agonist. *J. Med. Chem.* **1994**, *37*, 2828-2830.
 197. Glennon, R. A.; Herndon, J. L.; Dukat, M. Epibatidine-aided studies toward definition of a nicotine receptor pharmacophore. *Med. Chem. Res.* **1994**, *4*:461-473.
- 1995-1999**
198. Choudhary, M. S.; Sachs, N.; Ulver, A.; Glennon, R. A.; Westkaemper, R. B.; Roth, B. L. Differential ergoline and ergopeptide binding to 5-HT_{2A} receptors: Ergolines require an aromatic residue at position 340 for high affinity. *Mol. Pharmacol.* **1995**, *47*, 450-457.

199. Ismaiel, A. M.; Aruda, K.; Teitler, M.; Glennon, R. A. Ketanserin analogues: the effect of structural modification on 5-HT₂ serotonin receptor binding. *J. Med. Chem.* **1995**, *38*, 1196-1202.
200. Glennon, R. A.; Young, R.; Martin, B.; Dal Cason, T. Methcathinone ("CAT"): An enantiomeric potency comparison. *Pharmacol. Biochem.* **1995**, *50*, 601-601.
201. Hong, S.-S.; Dukat, M.; Teitler, M.; Herrick-Davis, K.; McCallum, K.; Kamboj, R.; Glennon, R. A. Binding of 5-arylalkoxytryptamines at human 5-HT_{1D8} serotonin receptors. *Med. Chem. Res.* **1995**, *5*, 690-699.
202. Glennon, R. A.; Hong, S.-S.; Bondarev, M.; Law, H.; Dukat, M.; Rakhit, S.; Power, P.; Fan, E.; Kinneau, D.; Kamboj, R.; Teitler, M.; Herrick-Davis, K.; Smith, C. Binding of O-alkyl derivatives of serotonin at human 5-HT_{1D8} receptors. *J. Med. Chem.* **1996**, *39*, 314-322.
203. Glennon, R. A.; Dukat, M.; Westkaemper, R. B.; Ismaiel, A. M.; Izzarelli, D. G.; Parker, E. M. The binding of propranolol at 5-hydroxytryptamine_{1D8} T355N mutant receptors may involve formation of two hydrogen bonds to asparagine. *Mol. Pharmacol.* **1996**, *49*, 198-206.
204. Young, R.; Glennon, R. A. A three-lever operant procedure differentiates the stimulus effects of R(-)-MDA from S(+)-MDA. *J. Pharmacol. Exp. Ther.* **1996**, *276*, 594-601.
205. Ismaiel, A. M.; Dukat, M.; Nelson, D. L.; Lucaites, V. L.; Glennon, R. A. Binding of N₂-substituted pyrido[4,3-b]indole analogs of spiperone at human 5-HT_{2A}, 5-HT_{2B}, and 5-HT_{2C} serotonin receptors. *Med. Chem. Res.* **1996**, *6*, 197-211.
206. Glennon, R. A.; Fiedler, W.; Dukat, M.; Damaj, I.; Martin, B. Pyrrolidine-modified analogs of nicotine: a structure-affinity investigation. *Eur. J. Med. Chem.* **1996**, *31*, 875-888.
207. Iwamura, T.; Casey, C. T.; Young, R.; Dukat, M.; Teitler, M.; Fadden, J. S. P.; Glennon, R. A. 4-(6-Fluorobenzisoxazol-3-yl)piperidine, a risperidone metabolite with serotonergic activity of potential clinical significance. *Med. Chem. Res.* **1996**, *6*, 593-601.
208. Glennon, R. A.; Dukat, M. Nicotine receptor ligands. *Med. Chem. Res.* **1996**, *6*, 465-486.
209. Dukat, M.; Abdel-Rahman, A. A.; Ismaiel, A. M.; Ingher, S.; Teitler, M.; Gyermek, L.; Glennon, R. A. Structure-activity relationships for the binding of arylpiperazines and arylbiguanides at 5-HT₃ serotonin receptors. *J. Med. Chem.* **1996**, *39*, 4017-4026.
210. Malmusi, L.; Dukat, M.; Young, R.; Teitler, M.; Darmani, N.; Ahmad, B.; Smith, C.; Glennon, R. A. 1,2,3,4-Tetrahydroisoquinolines and related analogs of the phenylalkylamine stimulants and hallucinogens. *Med. Chem. Res.* **1996**, *6*, 400-411.
211. Malmusi, L.; Dukat, M.; Young, R.; Teitler, M.; Darmani, N.; Ahmad, B.; Smith, C.; Glennon, R. A. 1,2,3,4-Tetrahydroisoquinolines and related analogs of the phenylalkylamine designer drug MDMA. *Med. Chem. Res.* **1996**, *6*, 412-426.
212. Damaj, M. I.; Glassco, W.; Dukat, M.; May, E. L.; Glennon, R. A.; Martin, B. R. Pharmacology of novel nicotinic analogs. *Drug Dev. Res.* **1996**, *38*, 177-187.
213. Young, R.; Darmani, N. A.; Elder, E. L.; Dumas, D.; Glennon, R. A. Clobenzorex: Evidence for amphetamine-like behavioral actions. *Pharmacol. Biochem. Behav.* **1997**, *56*, 311-316.
214. Glennon, R. A.; Dukat, M. Novel serotonergic agents: 5-HT₂ – Update 1997. *Investigational Drugs Research Alerts* **1997**, *2*, 107-113.
215. Glennon, R. A.; Young, R.; Dukat, M.; Cheng, Y. Initial characterization of PMMA as a discriminative stimulus. *Pharmacol. Biochem. Behav.* **1997**, *57*, 151-158.
216. Young, R. A.; Glennon, R. A. Cocaine-stimulus generalization to MDA optical isomers: A re-evaluation. *Pharmacol. Biochem. Behav.* **1997**, *57*, 115-118.
217. Glennon, R. A.; Dukat, M. 5-HT₁ receptor ligands: Update 1997. *Investigational Drugs Research Alerts* **1997**, *2*, 351-372.
218. Teitler, M.; Schieck, C.; Howard, P.; Sullivan, J. E.; Iwamura, T.; Glennon, R. A. 5-HT_{5A} serotonin receptors: A preliminary structure-affinity investigation. *Med. Chem. Res.* **1997**, *7*, 207-218.
219. Dal Cason, T.; Young, R.; Glennon, R. A. Cathinone: An investigation of several N-alkyl and methylenedioxy-substituted analogs. *Pharmacol. Biochem. Behav.* **1997**, *58*, 1109-1116.

220. Ismaiel, A. M.; Dukat, M.; Law, H.; Kamboj, R.; Lee, D. K. H.; Mazzocco, L.; Buekshkens, D.; Teitler, M.; Pierson, M. E.; Glennon, R. A. 2-(1-Naphthoxy)ethylamines with enhanced affinity for human 5-HT_{1DB} (h5-HT_{1B}) serotonin receptors. *J. Med. Chem.* **1997**, *40*, 4415-4419.
221. Grella, B.; Dukat, M.; Young, R.; Teitler, M.; Herrick-Davis, K.; Gauthier, C. B.; Glennon, R. A. Investigation of hallucinogenic and related β -carbolines. *Drug Alcohol Depend.* **1998**, *50*, 99-107.
222. Young, R.; Gabryszuk, M.; Glennon, R. A. (-)Ephedrine and caffeine mutually potentiate one another's amphetamine-like stimulus effects. *Pharmacol. Biochem. Behav.* **1998**, *61*, 169-173.
223. Young, R.; Glennon, R. A. Discriminative stimulus properties of (-)ephedrine. *Pharmacol. Biochem. Behav.* **1998**, *60*, 771-775.
224. Damaj, M. I.; Fei-Yin, M.; Dukat, M.; Glassco, W.; Glennon, R. A.; Martin, B. R. Antinociceptive responses to nicotinic acetylcholine receptor ligands after systemic and intrathecal administration in mice. *J. Pharmacol. Exp. Ther.* **1998**, *284*, 1058-1065.
225. Ablordeppey, S. Y.; Ashmawy, M.; Fischer, J. B.; Glennon, R. A. A CoMFA investigation of sigma receptor binding affinity: Reexamination of a spurious sigma ligand. *Eur. J. Med. Chem.* **1998**, *33*, 625-633.
226. Law, H.; Dukat, M.; Teitler, M.; Lee, D. K. H.; Mazzocco, L.; Kamboj, R.; Rampersad, V.; Prisinzano, T.; Glennon, R. A. Benzylimidazolines as h5-HT_{1B/1D} ligands: A structure-affinity investigation. *J. Med. Chem.* **1998**, *41*, 2243-2251.
227. Egan, C. T.; Herrick-Davis, K.; Miller, K.; Glennon, R. A.; Teitler, M. Agonist activity of LSD and lisuride at cloned 5HT_{2A} and 5HT_{2C} receptors. *Psychopharmacology* **1998**, *136*, 409-414.
228. Bondarev, M.; Iwamura, T.; Hynd, D.; Mazzocco, L.; Lee, D. K. H.; Dukat, M.; Glennon, R. A. Aryloxyethylamines as h5-HT_{1D} serotonin receptor ligands. *Med. Chem. Res.* **1998**, *8*, 333-342.
229. Metwally, K. A.; Dukat, M.; Egan, C. T.; Smith, C.; DuPre, A.; Gauthier, C. B.; Herrick-Davis, K.; Teitler, M.; Glennon, R. A. Spiperone: Influence of spiro ring substituents on 5-HT_{2A} serotonin receptor binding. *J. Med. Chem.* **1998**, *41*, 5084-5093.
230. Young, R.; Glennon, R. A. Discriminative stimulus effects of S(-)-methcathinone (CAT): A potent stimulant drug of abuse. *Psychopharmacology* **1998**, *140*, 250-256.
231. Dukat, M.; Dowd, M.; Damaj, M. I.; Martin, B.; El-Zahabi, M. A.; Glennon, R. A. Synthesis, receptor binding, and QSAR studies on 6-substituted nicotine derivatives as cholinergic ligands. *Eur. J. Med. Chem.* **1999**, *34*, 31-40.
232. Cheng, Y.-X.; Dukat, M.; Dowd, M.; Fiedler, W.; Martin, B.; Damaj, M. I.; Glennon, R. A. Synthesis and binding of 6,7,8,9-tetrahydro-5H-pyrido[3,4-d]azepine and related ring-opened analogs at central nicotinic receptors. *Eur. J. Med. Chem.* **1999**, *34*, 177-190.
233. Glennon, R. A.; Bondarev, M.; Roth, B. 5-HT₆ serotonin receptor binding of indolealkylamines: A preliminary structure-affinity investigation. *Med. Chem. Res.* **1999**, *9*, 108-117.
234. Nelson, D. L.; Lucaites, V. L.; Wainscott, D. B.; Glennon, R. A. Comparisons of hallucinogenic phenylisopropylamine binding affinities at cloned human 5-HT_{2A}, 5-HT_{2B}, and 5-HT_{2C} receptors. *Naunyn-Schmiedeberg's Arch. Pharmacol.* **1999**, *359*, 1-6.
235. Westkaemper, R. B.; Hyde, E. G.; Choudhary, M. S.; Khan, N.; Gelbar, E.; Glennon, R. A.; Roth, B. L. Engineering a region of bulk tolerance in the 5-HT_{2A} receptor. *Eur. J. Med. Chem.* **1999**, *34*, 441-447.
236. Young, R.; Bondarev, M.; Glennon, R. A. An examination of isomeric phenylpropanolamines in (-)ephedrine-trained rats. *Drug Alcohol Depend.* **1999**, *57*, 1-6.
237. Flammia, D.; Dukat, M.; Damaj, M. I.; Martin, B.; Glennon, R. A. Lobeline: A structure-affinity investigation of nicotinic acetylcholinergic (nACh) receptor binding. *J. Med. Chem.* **1999**, *42*, 3726-3731.
238. Westkaemper, R. B.; Runyon, S. P.; Bondarev, M. L.; Savage, J. E.; Roth, B.; Glennon, R. A. 9-(Aminomethyl)-9,10-dihydroanthracene is a novel and unlikely 5-HT_{2A} receptor antagonist. *Eur. J. Pharmacol.* **1999**, *380*, R5-R7.
239. Glennon, R. A. Arylalkylamine drugs of abuse: An overview of drug discrimination studies. *Pharmacol. Biochem. Behav.* **1999**, *64*, 251-256.

240. Khorana, N.; Bondarev, M.; Dukat, M.; Herrick-Davis, K.; Egan, C.; DuPre, A.; Smith, C.; Teitler, M.; Glennon, R. A. Binding of spiperone analogs at 5-HT_{2A} serotonin receptors. *Med. Chem. Res.* **1999**, *9*, 657-667.
 241. Young, R.; Dukat, M.; Malmusi, L.; Glennon, R. A. Stimulus properties of PMMA: Effect of optical isomers and conformational restriction. *Pharmacol. Biochem. Behav.* **1999**, *64*, 449-453.
 242. Hong, S.; Dukat, M.; Teitler, M.; Egan, C.; Dupre, A.; Herrick-Davis, K.; Glennon, R. A. 5,8-Dimethoxyharmalan: Actions at 5-HT_{2A} serotonin receptors. *Med. Chem. Res.* **1999**, *9*, 374-388.
- 2000-2004**
243. Glennon, R. A.; Lee, M.; Rangisetty, J. B.; Dukat, M.; Roth, B. L.; Savage, J. E.; McBride, A.; Rauser, L.; Hufeisen, S.; Lee, D. K. H. 2-Substituted tryptamines: Agents with selectivity for 5-HT₆ receptors. *J. Med. Chem.* **2000**, *43*, 1011-1018.
 244. Glennon, R. A.; Young, R. (+)Amphetamine stimulus generalization to an herbal ephedrine product. *Pharmacol. Biochem. Behav.* **2000**, *65*, 655-658.
 245. Kristiansen, K.; Kroeze, W. K.; Willins, D. L.; Gelber, E. I.; Savage, J. E.; Glennon, R. A.; Roth, B. L. A highly conserved aspartic acid (Asp-155) anchors the terminal amine moiety of tryptamines and is involved in membrane targeting of the 5-HT_{2A} serotonin receptor but does not participate in activation via a "Salt-bridge Disruption" mechanism. *J. Pharmacol. Exp. Ther.* **2000**, *293*, 735-746.
 246. Glennon, R. A.; Dukat, M. Central nicotinic receptor ligands and pharmacophores. *Pharm. Acta Helv.* **2000**, *74*, 103-114. Reprinted in book form; see entry 41 in the Chapter section below.
 247. Glennon, R. A.; Dukat, M.; Grella, B.; Hong, S.; Costantino, L.; Teitler, M.; Smith, C.; Egan, C.; Davis, K.; Mattson, M. V. Binding of β -carbolines and related agents at serotonin (5-HT₂ and 5-HT_{1A}), dopamine (D₂) and benzodiazepine receptors. *Drug Alcohol Depend.* **2000**, *60*, 121-132.
 248. Dowd, C. S.; Herrick-Davis, K.; Egan, C.; DuPre, A.; Teitler, M.; Glennon, R. A. 1-[4-(3-Phenylalkyl)phenyl]-2-aminopropanes as 5-HT_{2A} partial agonists. *J. Med. Chem.* **2000**, *43*, 3074-3084.
 249. Glennon, R. A.; Young, R. MDMA stimulus generalization to the 5-HT_{1A} serotonin agonist 8-OH DPAT. *Pharmacol. Biochem. Behav.* **2000**, *66*, 483-488.
 250. Young, R.; Glennon, R. A. Stimulus effects of phenylpropanolamine optical isomers in (+)amphetamine-trained rats. *Pharmacol. Biochem. Behav.* **2000**, *66*, 489-494.
 251. Dukat, M.; Young, R.; Darmani, N.; Ahmed, B.; Glennon, R. A. The 5-HT₃ agent N-(3-chlorophenyl)guanidine (MD-354) serves as a discriminative stimulus in rats and displays partial agonist character in a shrew emesis assay. *Psychopharmacology* **2000**, *150*, 200-207.
 252. Hellberg, M.; Stubbins, J. F.; Glennon, R. A. A preliminary investigation of mesoionic xanthine analogs as inhibitors of platelet aggregation. *Bioorg. Med. Chem. Lett.* **2000**, *8*, 1917-1923.
 253. Ablordepey, S. Y.; Fischer, J. B.; Glennon, R. A. Is a nitrogen atom an important pharmacophoric element in sigma ligand binding? *Bioorg. Med. Chem.* **2000**, *8*, 2105-2111.
 254. Tsai, Y.; Dukat, M.; Slassi, M.; MacLean, N.; Demchysyn, L.; Savage, J. E.; Roth, B. L.; Hufeisen, S.; Glennon, R. A. N₁-Benzenesulfonamidotryptamines as novel 5-HT₆ antagonists. *Bioorg. Med. Chem. Lett.* **2000**, *10*, 2295-2299.
 255. Ferretti, G.; Dukat, M.; Giannella, M.; Piergentili, A.; Pigini, M.; Quaglia, W.; Damaj, M. I.; Martin, B. R.; Glennon, R. A. Chain-lengthened and imidazoline analogs of nicotine. *Bioorg. Med. Chem. Lett.* **2000**, *10*, 2665-2668.
 256. Lee, M.; Rangisetty, J. B.; Dukat, M.; Slassi, A.; Maclean, N.; Lee, D. K. H.; Glennon, R. A. 5-HT₆ Serotonin receptor binding affinities of N₁-benzenesulfonyl and related tryptamines. *Med. Chem. Res.* **2000**, *10*, 230-242.
 257. Prinszanao, T.; Law, H.; Dukat, M.; Slassi, A.; Maclean, N.; Demchysyn, L.; Glennon, R. A. Imidazoline-modified benzylimidazolines as h5-HT_{1D/1B} serotonergic ligands. *Bioorg. Med. Chem.* **2001**, *9*, 613-619.

258. Westkaemper, R. B.; Runyon, S. P.; Savage, J. E.; Roth, B. L.; Glennon, R. A. Exploring the relationship between binding modes of 9-(aminomethyl)-9,10-dihydroanthracene and cyproheptadine analogs at the 5-HT_{2A} serotonin receptor. *Bioorg. Med. Chem. Lett.* **2001**, *11*, 563-566.
259. Runyon, S. P.; Savage, J. E.; Taroua, M.; Roth, B. L.; Glennon, R. A.; Westkaemper, R. B. Influence of chain length and N-alkylation on the selective serotonin receptor ligand 9-(aminomethyl)-9,10-dihydroanthracene. *Bioorg. Med. Chem. Lett.* **2001**, *11*, 655-658.
260. Prisinzano, T. P.; Law, H.; Slassi, A.; McClean, N.; Demchysyn, L.; Glennon, R. A. QSAR studies on 2-benzylimidazoline analogs as h5-HT_{1D/1B} serotonin receptor ligands. *Med. Chem. Res.* **2001**, *10*, 309-317.
261. Rangisetty, J. B.; Bondarev, M. L.; Chang-Fong, J.; Young, R.; Glennon, R. A. PMMA-Stimulus generalization to the optical isomers of MBDB and 3,4-DMA. *Pharmacol. Biochem. Behav.* **2001**, *69*, 261-267.
262. Dukat, M.; Choi, Y.; Teitler, M.; Du Pre, A.; Herrick-Davis, K.; Smith, C.; Glennon, R. A. The binding of arylguanidines at 5-HT₂ serotonin receptors: A structure-affinity investigation. *Bioorg. Med. Chem. Lett.* **2001**, *11*, 1599-1603.
263. Rangisetty, J. B.; Dukat, M.; Dowd, C. S.; Herrick-Davis, K.; Du Pre, A.; Gadepalli, S.; Teitler, M.; Kelley, C. R.; Sharif, N.; Glennon, R. A. 1-[2-Methoxy-5-(3-phenylpropyl)]-2-aminopropane unexpectedly shows 5-HT_{2A} serotonin receptor affinity and antagonist character. *J. Med. Chem.* **2001**, *44*, 3283-3291.
264. Hong, S.; Young, R.; Glennon, R. A. Discriminative stimulus properties of α -ethyltryptamine optical isomers. *Pharmacol. Biochem. Behav.* **2001**, *70*, 311-316.
265. Husbands, S. M.; Glennon, R. A.; Gogerat, S.; Gough, R.; Tyacke, R.; Crosby, J.; Nutt, D. J.; Lewis, J. W.; Hudson, A. L. β -Carboline binding to imidazoline receptors. *Drug Alcohol Depend.* **2001**, *64*, 203-208.
266. Chang-Fong, J.; Addo, J.; Dukat, M.; Smith, C.; Mitchell, N. A.; Herrick-Davis, K.; Teitler, M.; Glennon, R. A. Evaluation of isortryptamine derivatives at 5-HT₂ serotonin receptors. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 155-158.
267. Dukat, M.; Damaj, M. I.; Young, R.; Vann, R.; Collins, A. C.; Marks, M. J.; Martin, B. R.; Glennon, R. A. Functional diversity among 5-substituted nicotine analogs; *in vitro* and *in vivo* investigations. *Eur. J. Pharmacol.* **2002**, *435*, 171-180.
268. Young, R.; Glennon, R. A. The stimulus effects of 5,6,7,8-tetrahydro-1,3-dioxolo[4,5-g]isoquinoline is similar to that of cocaine but different from that of amphetamine. *Pharmacol. Biochem. Behav.* **2002**, *71*, 205-213.
269. Dukat, M.; Young, R.; Glennon, R. A. Effect of PMA optical isomers and 4-MTA in PMMA-trained rats. *Pharmacol. Biochem. Behav.* **2002**, *72*, 299-305.
270. Glennon, R. A.; Young, R. Effect of 1-(3,4-methylenedioxypheyl)-2-aminopropane and its optical isomers in PMMA-trained rats. *Pharmacol. Biochem. Behav.* **2002**, *72*, 307-311.
271. Glennon, R. A.; Young, R.; Rangisetty, J. B. Further characterization of the stimulus properties of 5,6,7,8-tetrahydro-1,3-dioxolo[4,5-g]isoquinoline. *Pharmacol. Biochem. Behav.* **2002**, *72*, 379-387.
272. Runyon, S. P.; Peddi, S.; Savage, J. E.; Roth, B. L.; Glennon, R. A.; Westkaemper, R. B. Geometry-affinity relationships of the selective serotonin receptor ligand 9-(aminomethyl)-9,10-dihydroanthracene. *J. Med. Chem.* **2002**, *45*, 1656-1664.
273. Glennon, R. A.; Metwally, K.; Dukat, M.; Ismaiel, A. M.; De Los Angeles, J.; Herndon, J.; Teitler, M.; Khorana, N. Ketanserin and spiperone as templates for novel serotonin 5-HT_{2A} antagonists. *Curr. Top. Med. Chem.* **2002**, *2*, 539-558.
274. Young, R.; Glennon, R. A. Nicotine and bupropion share a similar discriminative stimulus effect. *Eur. J. Pharmacol.* **2002**, *443*, 113-118.
275. Westkaemper, R. B.; Glennon, R. A. Application of ligand SAR, receptor modeling and receptor mutagenesis to the discovery and development of a new class of 5-HT_{2A} ligands. *Curr. Top. Med. Chem.* **2002**, *2*, 575-598.

276. Ablordeppey, S. Y.; Fischer, J. B.; Law, H.; Glennon, R. A. Probing the proposed phenyl-A region of the sigma-1 receptor. *Bioorg Med Chem.* **2002**, *10*, 2759-2765.
277. Lee, M.; Dukat, M.; Liao, L.; Flammia, D.; Damaj, I. M.; Martin, B.; Glennon, R. A. A comparison of the binding of three series of nicotinic ligands. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 1989-1992.
278. Dukat, M.; El-Zahabi, M.; Ferretti, G.; Damaj, M. I.; Marin, B. R.; Glennon, R. A. (-)-6-n-Propylnicotine antagonizes the antinociceptive effects of (-)-nicotine. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 3005-3007.
279. Ferretti, G.; Dukat, M.; Giannella, M.; Piergentili, A.; Pigini, M.; Quaglia, W.; Damaj, M. I.; Martin, B. R.; Glennon, R. A. Homoazanicotine: A structure-affinity study for nicotinic acetylcholinergic (nACh) receptor binding. *J. Med. Chem.* **2002**, *45*, 4721-4731.
280. Bondareva, T.; Young, R.; Glennon, R. A. Central stimulants as discriminative stimuli: Asymmetric generalization between (-)-ephedrine and S(+)-methamphetamine. *Pharmacol. Biochem. Behav.* **2002**, *74*, 157-162.
281. Young, R. A.; Glennon, R. A. Nicotine and bupropion share a similar discriminative stimulus effect. *Eur. J. Pharmacol.* **2002**, *443*, 113-118.
282. Ablordeppey, S. Y.; Fischer, J. B.; Law, H.; Glennon, R. A. Probing the proposed phenyl-A region of the sigma-1 receptor. *Bioorg Med Chem.* **2002**, *10*, 2759-2765.
283. Lee, M.; Dukat, M.; Liao, L.; Flammia, D.; Damaj, I. M.; Martin, B.; Glennon, R. A. A comparison of the binding of three series of nicotinic ligands. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 1989-1992.
284. Dukat, M.; El-Zahabi, M.; Ferretti, G.; Damaj, M. I.; Marin, B. R.; Glennon, R. A. (-)-6-n-Propylnicotine antagonizes the antinociceptive effects of (-)-nicotine. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 3005-3007.
285. Ferretti, G.; Dukat, M.; Giannella, M.; Piergentili, A.; Pigini, M.; Quaglia, W.; Damaj, M. I.; Martin, B. R.; Glennon, R. A. Homoazanicotine: A structure-affinity study for nicotinic acetylcholinergic (nACh) receptor binding. *J. Med. Chem.* **2002**, *45*, 4721-4731.
286. Bondareva, T.; Young, R.; Glennon, R. A. Central stimulants as discriminative stimuli: Asymmetric generalization between (-)-ephedrine and S(+)-methamphetamine. *Pharmacol. Biochem. Behav.* **2002**, *74*, 157-162.
287. Young, R. A.; Glennon, R. A. Nicotine and bupropion share a similar discriminative stimulus effect. *Eur. J. Pharmacol.* **2002**, *443*, 113-118.
288. Ferretti, G.; Dukat, M.; Giannella, M.; Piergentili, A.; Pigini, M.; Quaglia, W.; Damaj, M. I.; Martin, B. R.; Glennon, R. A. Binding of nicotine and homoazanicotine at neuronal nicotinic acetylcholine (nACh) receptors. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 733-735.
289. Rahman, A. A.; Daoud, M. K.; Dukat, M.; Herrick-Davis, K.; Purohit, A.; Teitler, M.; Taveres do Ameral, A.; Malvezzi, A.; Glennon, R. A. Conformationally-restricted analogues and partition coefficients of the 5-HT₃ serotonin receptor ligands meta-chlorophenylbiguanide (mCPBG) and meta-chlorophenylguanidine (mCPG). *Bioorg. Med. Chem. Lett.* **2003**, *13*, 1119-1123.
290. Khorana, N.; Purohit, A.; Herrick-Davis, K.; Teitler, M.; Glennon, R. A. γ -Carbolines: Binding at 5-HT_{3A} serotonin receptors. *Bioorg. Med. Chem.* **2003**, *11*, 717-722.
291. Dukat, M.; Glennon, R. A. Epibatidine: Impact on nicotine receptor research. *Cell. Molec. Neurobiol.* **2003**, *23*, 365-378.
292. Setola, V.; Hufeisen, S. J.; Grande-Allen, K. J.; Vesely, I.; Glennon, R. A.; Blough, B.; Rothman, R. B.; Roth, B. L. 3,4-Methylenedioxymethamphetamine (MDMA, "Ecstasy") induces fenfluramine-like proliferative actions on human cardiac valvular interstitial cells in vitro. *Mol. Pharmacol.* **2003**, *63*, 1223-1229.
293. Glennon, R. A. Higher-end serotonin receptors: 5-HT₅, 5-HT₆, and 5-HT₇. *J. Med. Chem.* **2003**, *46*, 2795-2812.
294. Peddi, S.; Roth, B. L.; Glennon, R. A.; Westkaemper, R. B. Ring substituted analogues of 5-aminomethyl-10,11-dihydrobenzo[a,d]cycloheptene (AMDA): potential modes of binding to the 5-HT_{2A} receptor. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 2565-2568.

295. Eckler, J. R.; Chang-Fong, J.; Rabin, R. A.; Smith, C.; Teitler, M.; Glennon, R. A.; Winter, J. C. Behavioral characterization of 2-O-desmethyl and 5-O-desmethyl metabolites of the phenylethylamine hallucinogen DOM. *Pharmacol. Biochem. Behav.* **2003**, *75*, 845-852.
296. Simsek, R.; Chang-Fong, J.; Lee, M.; Damaj, M. I.; Martin, B. R.; Glennon, R. A. Quaternary ammonium 3-(aminoethoxy)pyridines as antinociceptive agents. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 2917-2920.
297. Bondarev, M. L.; Bondareva, T. S.; Young, R.; Glennon, R. A. Behavioral and biochemical investigations of bupropion metabolites. *Eur. J. Pharmacol.* **2003**, *474*, 85-93.
298. Pullagurla, M. R.; Dukat, M.; Setola, V.; Roth, B.; Glennon, R. A. N¹-Benzenesulfonylgramine and N¹-benzenesulfonylskatole: Novel 5-HT₆ receptor ligand templates. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 3355-3359.
299. Khorana, N.; Smith, C.; Herrick-Davis, K.; Purohit, A.; Teitler, M.; Grella, B.; Dukat, M.; Glennon, R. A. Binding of tetrahydro- γ -carbolines at human 5-HT_{2A} receptors. *J. Med. Chem.* **2003**, *46*, 3930-3937.
300. Rothman, R. B.; Vu, N.; Partilla, J. S.; Roth, B. L.; Young, R.; Glennon, R. A. In Vitro Characterization of echedrine-like phenylpropanolamine stereoisomers at biogenic amine transporters and the receptorsome reveals selective action as norepinephrine transporter substrates. *J. Pharmacol. Exp. Ther.* **2003**, *307*, 138-145.
301. Glennon, R. A.; Daoud, M. K.; Dukat, M.; Teitler, M.; Herrick-Davis, K.; Purohit, A.; Syed, H. Arylguanidine and arylbiguanide binding at 5-HT₃ serotonin receptors: A QSAR study. *Bioorg. Med. Chem. Lett.* **2003**, *11*, 4449-4454.
302. Glennon, R. A. Higher end Serotonin Receptors: 5-HT₅, 5-HT₆, 5-HT₇. *J. Med. Chem.* **2003**, *46*, 2795-2812.
303. Grella, B.; Teitler, M.; Smith, C.; Herrick-Davis, K.; Glennon, R. A. Binding of β -carbolines at 5-HT₂ serotonin receptors. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 4421-4425.
304. Peddi, S.; Roth, B. L.; Glennon, R. A.; Westkaemper R. B. Spiro[9,10-dihydroanthracene]-9,3'-pyrrolidine – a structurally unique tetracyclic 5-HT_{2A} receptor antagonist. *Eur. J. Pharmacol.* **2003**, *482*, 335-337.
305. Dogruer, D.; Lee, M.; Dukat, M.; Damaj, M. I.; Martin, B. R.; Glennon, R. A. 3-(4-Aminobutyn-1-yl)pyridines: Binding at α 4 β 2 nicotinic cholinergic receptors. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 523-526.
306. Glennon, R. A.; Grella, B.; Tyacke, R. J.; Lau, A.; Westaway, J.; Hudson, A. L. Binding of an imidazopyridindole at imidazoline I₂ receptors. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 527-529.
307. Peddi, S.; Roth, B. L.; Glennon, R. A.; Westkaemper R. B. Structural Determinants for high 5-HT_{2A} receptor affinity of spiro[9,10-dihydroanthracene]-9,3'-pyrrolidine (SpAMDA). *Bioorg. Med. Chem. Lett.* **2004**, *14*, 2279-83.
308. Glennon, R. A.; Dukat, M.; α 4 β 2 nACh Receptor Pharmacophore Models. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 1841-1844.
309. Glennon, R. A.; Dukat, M. Musings on α 4 β 2 nicotinic acetylcholine (nACh) receptor pharmacophore models. *Curr. Topics Med. Chem.* **2004**, *4*, 631-644.
310. Glennon, R. A.; Grella, B.; Tyacke, R. J.; Lau, A.; Westaway, J.; Hudson, A. L. Binding of β -carbolines at imidazoline I₂ receptors: Structure-affinity investigation. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 999-1002.
311. Chang-Fong, J.; Tyacke, R. J.; Lau, A.; Westaway, J.; Hudson, A. L.; Glennon, R. A. Pyrazino[1,2-a]indoles as novel high-affinity and selective imidazoline I₂ receptor ligands. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 1003-1005.
312. Tyacke, R. J.; Lau, A.; Grella, B.; Glennon, R. A.; Nutt, D. J.; Hudson, A. L. Investigation of the affinities of two new β -carbolines for rat brain imidazoline₂ receptors. *Ann. N. Y. Acad. Sci.* **2004**, *1009*, 361-363.
313. Chang-Fong, J.; Rangisetty, J. B.; Dukat, M.; Setola, V.; Raffay, T.; Roth, B.; Glennon, R. A. 1,2,3,4-Tetrahydrocarbazoles as 5-HT₆ serotonin receptor ligands. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 1961-1964.

314. Glennon, R. A.; Ismaiel, A. M.; Ablordeppey, S.; El-Ashmawy, M.; Fisher, J. B. Thioxanthene-derived analogs as sigma-1 receptor ligands. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 2217-2220.
 315. Dukat, M.; Smith, C.; Herrick-Davis, K.; Teitler, M.; Glennon, R. A. Binding of tryptamine analogs at *h5-HT_{1E}* receptors: A structure-affinity investigation. *Bioorg. Med. Chem.* **2004**, *12*, 2545-2552.
 316. Khorana, N.; Pullagurula, M. R.; Young, R.; Glennon, R. A. Comparison of the discriminative stimulus effects of 3,4-methylenedioxymethamphetamine (MDMA) and cocaine: Asymmetric generalization. *Drug Alcohol Depend.* **2004**, *74*, 281-287.
 317. Dukat, M.; Taroua, M.; Dahdouh, A.; Siripurapu, U.; Damaj, M.; Martin, B. R.; Glennon, R. A. (\pm)-8-Amino-5,6,7,8-tetrahydroisoquinolines as novel antinociceptive agents. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 3651-3654.
 318. Pullagurula, M. R.; Bondareva, T.; Young, R.; Glennon, R. A. Modulation of the stimulus effects of (+)amphetamine by the 5-HT₆ antagonist MS-245. *Pharmacol. Biochem. Behav.* **2004**, *78*, 263-268.
 319. Pullagurula, M. R.; Westkaemper, R. B.; Glennon, R. A. Possible differences in modes of binding of agonist and antagonist binding at human 5-HT₆ receptors. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 4569-4573.
 320. Prisinzano, T.; Dukat, M.; Law, H.; Slassi, A.; MacLean, N. A.; DeLannoy, I.; Glennon, R. A. 2-(Anilino)imidazolines and 2-(benzyl)imidazoline derivatives as *h5-HT_{1D}* serotonin receptor ligands. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 4697-4699.
 321. Khorana, N.; Pullagurula, M. R.; Dukat, M.; Young, R.; Glennon, R. A. Stimulus effects of three sulfur-containing psychoactive agents. *Pharmacol. Biochem. Behav.* **2004**, *78*, 821-826.
 322. Glennon, R. A. Nicotine and pain. *Med. Chem. Res.* **2004**, *13*, 74-77.
 323. Glennon, R. A.; Bondarev, M. L.; Khorana, N.; Young, R.; May, J. A.; Hellberg, R. R.; McLaughlin, M. A.; Sharif, N. A. β -Oxygenated analogs of the 5-HT_{2A} serotonin agonist 1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane (DOB). *J. Med. Chem.* **2004**, *47*, 6034-6041.
- 2005-**
324. Lee, M.; Rangisetty, J. B.; Pullagurula, M. R.; Dukat, M.; Setola, V.; Roth, B. L.; Glennon, R. A. 1-(1-Naphthyl) piperazines as a novel template for 5-HT₆ serotonin receptor ligands. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 1707-1711.
 325. Kolanos, R.; Siripurapu, U.; Pullagurula, M. R.; Riaz, M.; Setola, V.; Roth, B. L.; Glennon, R. A. Binding of iso- tryptamines and indenes at *h5-HT₆* receptors. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 1987-1991.
 326. Setola, V.; Dukat, M.; Glennon, R. A.; Roth, B. L. Molecular determinants for the interaction of the vavulopathic anorexigen norfenfluramine with the 5-HT_{2B} receptor. *Mol. Pharmacol.* **2005**, *68*, 20-33.
 327. Ramunno, A.; Dukat, M.; Lee, M.; Young, R.; El-Zahabi, M.; Damaj, M. I.; Martin, B.; Glennon, R. A. 6-(2-Phenylethyl)nicotine: A novel nicotinic cholinergic receptor ligand. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 3237- 3240.
 328. Abate, C.; Kolanos, R.; Dukat, M.; Setola, V.; Roth, B. L.; Glennon, R. A. Interaction of chiral MS-245 analogs at 5-HT₆ receptors. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 3510-3513.
 329. Dukat, M.; Ramunno, A.; Banzi, R.; Damaj, I. M.; Martin, B.; Glennon, R. A. 3-(2-Aminoethyl)pyridine analogs as $\alpha 4\beta 2$ nicotinic cholinergic receptor ligands. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 4308-4312.
 330. Glennon, R. A. Pharmacophore identification for σ_1 receptor binding. *Mini-Rev. Med. Chem.* **2005**, *5*, 927-940.
 331. Pullagurula, M.; Dukat, M.; Roth, B. L.; Setola, V.; Glennon, R. A. 5-Azatriptamine analogs as *h5-HT₆* receptor ligands. *Med. Chem. Res.* **2005**, *14*, 1-18.
 332. Pullagurula, M.; Siripurapu, U.; Kolanos, R.; Bondarev, M. L.; Dukat, M.; Setola, V.; Roth, B. L.; Glennon, R. A. Binding of amine-substituted N₁-benzenesulfonylindoles at human 5-HT₆ serotonin receptors. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 5298-5302.

333. Young, R.; Khorana, N.; Bondareva, T.; Glennon, R. A. Pizotyline effectively attenuates the stimulus effects of N-methyl-3,4-methylenedioxamphetamine (MDMA). *Pharmacol. Biochem. Behav.* **2005**, *82*, 404-410.
334. Glennon, R. A. Binding characteristics of σ_2 receptor ligands. *Brazil. J. Pharm. Sci.* **2005**, *41*, 1-12.
335. Bondareva, T.; Wesolowska, A.; Dukat, M.; Lee, M.; Young, R.; Glennon, R. A. S(+) and R(-)-N-methyl-1-(3,4-methylenedioxyphenyl)-2-aminopropane (MDMA) as discriminative stimuli: effect of cocaine. *Pharmacol. Biochem. Behav.* **2005**, *82*, 531-538.
336. Young, R.; Bondareva, T.; Wesolowska, A.; Glennon, R. A. Modulation of a (+)amphetamine discriminative stimulus in rats by 8-hydroxy-2-(N,N-di-n-propylamino)tetralin (8-OH DPAT). *Pharmacol. Biochem. Behav.* **2006**, *83*, 612-617.
337. Siripurapu, U.; Kolanos, R.; Dukat, M.; Roth, B. L.; Glennon, R. A. Binding of methoxy-substituted N₁-benzenesulfonylindole analogs at human 5-HT₆ serotonin receptors. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 3793-6.
338. Young, R.; Rothman, R. B.; Rangisetty, J. B.; Partilla, J. S.; Dukat, M.; Glennon, R. A. TDIQ (5,6,7,8-tetrahydro-1,3-dioxolo[4,5-g]isoquinoline) inhibits the consumption of "snacks" in mice. *Pharmacol. Biochem. Behav.* **2006**, *84*, 74-83.
339. Young, R.; Batkai, S.; Dukat, M.; Glennon, R. A. TDIQ (5,6,7,8-tetrahydro-1,3-dioxolo[4,5-g]isoquinoline) exhibits anxiolytic-like activity in a marble-burying assay in mice. *Pharmacol. Biochem. Behav.* **2006**, *84*, 62-73.
340. Ragab, H. M.; Kim, J. S.; Dukat, M.; Navarro, H.; Glennon, R. A. Aryloxyethylamines: binding at alpha7 nicotinic acetylcholine receptors. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 4283-4276.
341. Sikazwe, D.; Bondarev, M. L.; Dukat, M.; Rangisetty, J. B.; Roth, B. L.; Glennon, R. A. Binding of sulfonyl-containing arylalkylamines at human 5-HT₆ serotonin receptors. *J. Med. Chem.* **2006**, *49*, 5217-5225.
342. Kolanos, R.; Dukat, M.; Roth, B. L.; Glennon, R. A. Interaction of N₁-unsubstituted and N₁-benzenesulfonyltryptamines at h5-HT₆ receptors. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 5832-5835.
343. Young, R.; Bondareva, T.; Wesolowska, A.; Young, S.; Glennon, R. A. Effect of the 5-HT₆ serotonin antagonist MS-245 on the actions of (-)nicotine. *Pharmacol. Biochem. Behav.* **2006**, *83*, 612-617.
344. Glennon, R. A.; Bondareva, T.; Young, R. α -Ethyltryptamine (alpha-ET) as a discriminative stimulus in rats. *Pharmacol. Biochem. Behav.* **2006**, *85*, 448-453.
345. Glennon, R. A. 3-(4-Tetrahydropyridin-1-yl)butyl)oxindoles as 5-HT₇ receptor ligands. *Expert Opin. Ther. Patents* **2006**, *16*, 1171-1174.
346. Nyandenge, A.; Kolanos, R.; Roth, B. L.; Glennon, R. A. Further studies on the binding of N₁-substituted tryptamines at h5-HT₆ receptors. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 1691-1694.
347. Glennon, R. A.; Young, R.; Dukat, M.; Chang-Fong, J.; El-Zahabi, M. N-Methyl-1-(4-methoxyphenyl)-2-aminopropane (PMMA) and N-Methyl-1-(3,4-methylenedioxyphenyl)-2-aminopropane (MDMA) produce non-identical discriminative stimuli in rats. *Pharmacol. Biochem. Behav.* **2007**, *86*, 477-484.
348. Dukat, M.; Glennon, R. A.; Young, S. MD-354: what is it good for? *CNS Drug Rev.* **2007**, *13*, 1-20.
349. Dukat, M.; Wesolowska, A.; Young, R.; Glennon, R. A. The 5-HT₇ receptor partial agonist MD-354 (meta-chlorophenylguanidine) enhances the discriminative stimulus actions of (+)amphetamine in rats. *Pharmacol. Biochem. Behav.* **2007**, *87*, 203-207.
350. Kim, J. S.; Padnya, A.; Weltzin, M.; Edmonds, B. W.; Schulte, M. K.; Glennon, R. A. Synthesis of desformylflustrabromine and its evaluation as an alpha4beta2 and alpha7 nACh receptor modulator. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 4855-60.
351. Young, R.; Glennon, R. A. MDMA (N-methyl-3,4-methylenedioxamphetamine) and its stereoisomers: Similarities and differences in behavioral effects in an automated activity apparatus in mice. *Pharmacol. Biochem. Behav.* **2008**, *88*, 318-331.
352. Duakt, M.; Mosier, P.D.; Kolanos, R.; Roth, B.L.; Glennon, R.A. Binding of serotonin and N₁-benzenesulfonyltryptamine-related analogs at human 5-HT₆. *J. Med. Chem.* **2008**, *51*, 603-611.

INVITED BOOK CHAPTERS

1. Glennon, R. A.; Roscraens, J. A.; Young, R. The use of the drug discrimination paradigm for studying hallucinogenic agents. In: **Drug Discrimination: Applications in CNS Pharmacology**, F. C. Colpaert and J. L. Slangen eds., Elsevier Biomedical Press, Amsterdam, 1982. pp 69-96.
2. Glennon, R. A. Hallucinogenic Phenylisopropylamines: Stereochemical Aspects. In: **Stereoisomers: Drugs in Psychopharmacology**, D. Smith ed., CRC Press, Boca Raton, 1983. pp 327-370.
3. Nichols, D. E.; Glennon, R. A. Medicinal Chemistry and Structure-Activity Relationships of Hallucinogenic Agents. In: **Hallucinogens: Neurochemical, Behavioral, and Clinical Perspectives**, B. Jacobs ed., Raven Press, NY 1984. pp 95-142.
4. Glennon, R. A. Interaction of hallucinogenic agents with serotonin receptors. In: **Medicinal Chemistry**, R. Dahldom and J. L. G. Nilsson, eds., Swedish Pharmaceutical Press, Stockholm 1985. pp 116-129.
5. Glennon, R. A. The role of serotonin in the mechanism of action of hallucinogenic agents. In: **Neuropharmacology of Serotonin**, A. R. Green, ed., Oxford University Press, 1985.
6. Arvidsson, L. E.; Hacksell, U.; Glennon, R. A. Recent advances in central 5-hydroxytryptamine receptor agonists and antagonists. In: **Progress in Drug Research**, E. Jucker, ed., Birkhauser Verlag, Basel, 1986. pp 365-471.
7. Glennon, R. A. Discriminative stimulus properties of site-selective serotonin agonists. In: **Transduction Mechanisms of Drug Stimuli**, F. C. Colpaert and R. L. Balster eds., Springer Verlag, Berlin, 1988. pp 15-32.
8. Glennon, R. A.; Young, R. Use of the drug discrimination paradigm for investigating structure-activity relationships, in **Drug Dependence**, M. Bozarth ed., Springer-Verlag, NY, 1987. pp 373-390.
9. Glennon, R. A. Psychoactive Phenylisopropylamines. In: **Psychopharmacology, Third Generation of Progress**, H. Meltzer ed., Raven Press, 1987. pp 1627-1634.
10. Young, R.; Glennon, R. A. Second generation anxiolytics and serotonin, In: **Serotonin: Behavioral Pharmacology**, R. H. Rech and G. A. Gudelsky eds. NPP Press, Ann Arbor MI, 1988. pp 239-258.
11. Glennon, R. A.; Lucki, I. Behavioral aspects of serotonin agonists", In: **Serotonin Receptors**, E. Sanders-Bush ed., Humana Press, NJ, 1988. pp 253-294.
12. Glennon, R. A. Central serotonin receptors, In: **Receptor Pharmacology and Function**, M. Williams, R. A. Glennon and P. Timmermans eds., M. Dekker, 1989. pp 257-292.
13. Glennon, R. A.; Young, R.; Pierson, M. E. Stimulus properties of arylpiperazine second generation anxiolytics. In: **The Behavioral Pharmacology of 5-HT**, P. Bevan, A. R. Cools, and T. Archer, eds. Lawrence Earlbaum Associates, NY, 1989. pp 445-449.
14. Glennon, R. A.; Seggel, M. R. Interaction of phenylisopropylamines with 5-HT₂ receptors: A QSAR analysis. In: **Bioactive Mechanisms**, Magee, P. S. ed. ACS Press, Washington DC, 1989.
15. Glennon, R. A. Synthesis and evaluation of amphetamine analogs: A review. In: **Clandestinely Produced Drugs, Analogues, and Precursors**, M. Klein, F. Sapienza, H. McClain and I. Khan, eds. US Government Printing Office, Washington DC, 1989. pp 39-65.
16. Glennon, R. A. "Stimulus Properties of Hallucinogenic Phenalkylamines and Related Designer Drugs: Formulation of Structure-Activity Relationships" In: **Pharmacology and Toxicology of Amphetamine and Related Designer Drugs**, Ashgar, R. DeSouza, E. eds., U.S. Government Printing Office, Washington, DC., 1989 pp 43-67.
17. Teitler, M.; Leonhardt, S.; Appel, N. M.; DeSouza, E. B.; Glennon, R. A. "Receptor pharmacology of MDMA and related hallucinogens" In: **The Neuropharmacology of Serotonin**, Peroutka, S., ed., New York Academy of Science, New York, 1990. pp. 626-638.
18. Glennon, R. A., Westkaemper, R., Bartyzel, P. "Medicinal Chemistry of Serotonergic Agents" In: **Serotonin Receptor Subtypes**, Peroutka, S. ed., Wiley-Liss, NY, 1991. pp. 19-64.
19. Glennon, R. A. Serotonergic agents and CNS receptors, in **Advances in CNS Drug-Receptor Interactions**, J. Cannon, ed. JAI Press, Greenwich. 1991.
20. Glennon, R. A., Tejani, S. "Mesoionic Nucleosides and Heterobases" In: **Chemistry of Nucleosides and Nucleotides**, L. Townsend, ed., Plenum Press, 1991. pp. 1-21.

21. Glennon, R.A., Peroutka, S.J., Dukat, M. "Binding characteristics of a quaternary amine analog of serotonin: 5-HTQ", In: **Serotonin: Molecular Biology, Receptors and Functional Effects**, J. R. Fozard and P. R. Saxena, eds., Birkhauser-Verlag, Basel, 1991. pp. 186-191.
22. Glennon, R. A. "Animal Models for Assessing Hallucinogenic Agents" In: **Animal Models for Assessing Hallucinogenic Agents**, Boulton, A.; Baker, G.; Wu, P.; ed., Humana Press, Clifton, New Jersey, 1992. pp. 345-381.
23. Glennon, R.A. "Discriminative Stimulus Properties of Hallucinogens and Related Designer Drugs" In: **Drug Discrimination: Application in Drug Abuse Research**, NIDA Research Monograph, R.A. Glennon, T. Jarbe and J. Frankenhein, eds., Washington DC, 1991. pp. 25-44.
24. Glennon, R.A. "Phenalkylamine Stimulants, Hallucinogens and Designer Drugs" In: **Proceedings of the 1990 CPDD Meeting**, L.S. Harris, ed., Washington DC, 1990. pp. 154-160.
25. Glennon, R.A.; Westkaemper, R.B. "Serotonin Receptors, 5-HT Ligands and Receptor Modeling" In: **Trends in Receptor Research**, P. Angeli, U. Gulini and W. Quaglia, eds, Elsevier, Amsterdam, 1992. pp. 185-207.
26. Glennon, R.A. Classification of agents used in serotonin research. In: **Serotonin**, P. M. Vanhoutte, P. R. Saxena, R. Paoletti, N. Brunello, A. S. Jackson, eds., Kluwer Academic Publishers, Dordrecht, 1993. pp 49-54.
27. Glennon, R.A. "Classical hallucinogens, an introductory overview". In: **Hallucinogens: An Update**, G.C. Lin and R.A. Glennon, eds., U.S. Government Printing Office, Washington, D.C. 1994. pp. 4-32.
28. Westkaemper, R.B.; Glennon, R.A. "Molecular modeling of the interaction of LSD and other hallucinogens with 5-HT₂ receptors". In: **Hallucinogens: An Update**, G. C. Lin and R.A. Glennon, eds., U. S. Government Printing Office, Washington, D.C. 1994. pp. 263-283.
29. Herndon, J.L.; Glennon, R.A. Serotonin receptors, agents, and actions. In: **Drug Design for Neuroscience**, A.P. Kozikowski, ed. Raven Press, NY, 1993. pp 167-212.
30. Glennon, R.A.; Dukat, M. "Serotonin Receptor Subtypes". In: **Psychopharmacology: The fourth generation of progress**, Bloom, R.E.; Kupfer, D.J.; Raven Press 1994, 415-429.
31. Glennon, R.A. How do LSD and LSD-related hallucinogens work? In: **Hallucinogens, LSD, and Raves**. Proceedings of the Scientific Advisory Board of the American Council for Drug Education, American Council for Drug Education, Washington, D. C. 1994. pp 14-20.
32. Glennon, R.A.; Dukat, M. Serotonin Subfamily of receptors. In: **The RBI Handbook of Receptor Classification**, J.W. Keabian and J.L. Neumeyer, eds., Research Biochemicals International, Natick, 1994. pp. 58-61..
33. Glennon, R.A. Sigma receptors. In: **The RBI Handbook of Receptor Classification**, J. W. Keabian and J. L. Neumeyer, eds., Research Biochemicals International, Natick, 1994. pp 62-63.
34. Dukat, M.; Damaj, M.I.; Dumas, D.; Glassco, W.; May, E.L.; Martin, B.R.; Glennon, R.A. "Epibatidine". In: **Effects of Nicotine on biological systems II (APS)**, P.B.S. Clarke, M. Quik, K. Thurau, F. Adlkofer, eds., Birkhauser, Basel, Switzerland, 1995.
35. Glennon, R.A. "Classical Hallucinogens". In: **Handbook of Experimental Pharmacology: Pharmacological Aspects of Drug Dependence**, C. R. Schuster and M. J. Kuhar, eds., Springer Verlag, Basel, 1996. pp 343-371.
36. Glennon, R.A.; Dukat, M. Serotonin receptors and ligands. In: **Introduction to the Principles of Drug Design and Action**, H. J. Smith, ed. 1997.
37. Glennon, R. A.; Dukat, M. Nicotinic receptor pharmacophores. In: **Neuronal Nicotinic Receptors: Pharmacology and Therapeutic Opportunities**, S. P. Arneric and J. D. Brioni, eds., John Wiley and Sons, N.Y., 1999, pp 271-284.
38. Glennon, R. A.; Dukat, M.; Westkaemper, R. B. Serotonin receptor subtypes and ligands. **Psychopharmacology, A Generation of Progress**, 1999, CD ROM Version. [See also: <http://www.acnp.org/citations/GN401000039/Default.htm>.]
39. Glennon, R. A. Pharmacology of hallucinogens, In: **Handbook of Substance Abuse: Neurobehavioral Pharmacology**, R. E. Tartar, R. T. Ammerman, and P. J. Ott, eds, Plenum Press, N.Y., 1999, pp 217-227.

40. Glennon, R. A. Neurobiology of hallucinogens, In: Textbook of Substance Abuse Treatment, M. Galanter and H. D. Kleber, eds., American Psychiatric Press, Washington, D. C., 1999, pp 33-37.
40. Glennon, R. A.; Dukat, M. Nicotine analogs: Structure-affinity relationships for central acetylcholinergic receptor binding. In: Nicotinic Agents, I. Yamamoto and J. Casida, eds., Springer-Verlag, Tokyo, 1999, pp 237-252.
41. Glennon, R. A.; Dukat, M. Central nicotinic receptor ligands and pharmacophores. In: Receptor Chemistry: Toward the Third Millennium, U. Gulini, M. Giannella, W. Quaglia, G. Marucci, eds., Elsevier, Amsterdam, 2000, pp 103-114.
42. Glennon, R. A. Hallucinogens, stimulants, and related drugs of abuse, In: Fove's Textbook of Medicinal Chemistry, D. A. Williams and T. Lemke, eds., Williams and Wilkins, Baltimore, 2002, pp 434-452.
43. Glennon, R. A.; Dukat, M. Serotonin receptors and drugs affecting serotonergic neurotransmission, In: Fove's Textbook of Medicinal Chemistry, D. A. Williams and T. Lemke, eds., Williams and Wilkins, Baltimore, 2002, pp 315-337.
44. Glennon, R. A. The pharmacology of serotonergic hallucinogens and designer drugs. In: Principles of Addiction Medicine, Third Edition, A. W. Graham, T. K. Schultz, M. Mayo-Smith, R. K. Ries and B. B. Wilford, eds., American Society of Addiction Medicine, Chevy Chase, MD. 2003, pp 275-281.
45. Iverson, L and Glennon, R. A. Antidepressants, In: Burger's Medicinal Chemistry and Drug Discovery, Volume 6, Nervous System Agents, D. J. Abraham, ed., Wiley: New York. 2003. Chapter 8, pgs 483-524.
46. Glennon, R. A. Medicinal chemistry of $\alpha 4\beta 2$ nicotinic cholinergic receptor ligands, In: Progress in Medicinal Chemistry, Volume 42, Elsevier Science B. V., Amsterdam, 2004, 55-123.
47. Glennon, R. A. Neurobiology of hallucinogens. In: Textbook of Substance Abuse Treatment, M. Galanter and H. D. Kleber, eds., American Psychiatric Publishing, Washington DC, 2004, pp 41-46.
48. Ablordeppey, S. Y.; Glennon, R. A. Pharmacophore models for sigma₁ receptor binding, In: Sigma Receptors, Chemistry, Cell Biology and Clinical Implications, Matsumoto, Rae, R.; Bowen, Wayne, D.; Su, Tsung-Ping (Eds.), 2007.
49. Glennon, R. A. Strategies for the development of selective serotonergic agents. In: The Serotonin Receptors: From Molecular Pharmacology to Human Therapeutics, B. L. Roth, ed., Humana Press, NJ., *in press*.
50. Glennon, R. A. Hallucinogens, stimulants, and related drugs of abuse, In: Fove's Textbook of Medicinal Chemistry, D. A. Williams and T. Lemke, eds., Williams and Wilkins, Baltimore, *in press*.
51. Glennon, R. A.; Dukat, M. Serotonin receptors and drugs affecting serotonergic neurotransmission, In: Fove's Textbook of Medicinal Chemistry, D. A. Williams and T. Lemke, eds., Williams and Wilkins, Baltimore, *in press*.
52. Glennon, R. A. Neurobiology of hallucinogens. In: Textbook of Substance Abuse Treatment, M. Galanter and H. D. Kleber, eds., American Psychiatric Publishing, Washington DC, *in press*.

BOOKS EDITED

1. **Receptor Pharmacology and Function**. M. Williams, R. A. Glennon and P. Timmermans eds., M. Dekker, New York, 1989.
2. **Drug Discrimination: Applications in Drug Abuse Research**, R. A. Glennon, T. Jarbe and J. Frankenheim, U.S. Government Printing Office, Washington DC, 1991.
3. **Hallucinogens: An Update**, G.C. Lin and R.A. Glennon, eds, U.S. Government Printing Office, Washington, D.C. 1994.

Patents

- Selective serotonin receptor antagonists and therapeutic applications thereof. United States Patent 6,806,283, October 19, 2004 (with R. B. Westkaemper)
- Selective 5-HT₆ receptor ligands. United States Patent 6,518,297, February 11, 2003
- Selective 5-HT₆ receptor ligand United States Patent 6,489,488, December 3, 2002
- Selective 5-HT₆ receptor ligands, United States Patent 6,403,808, June 11, 2002
- Imidazoles with serotonin receptor binding activity, United States Patent 6,288,101, September 11, 2001
- Imidazoles with serotonin receptor binding activity, United States Patent 6,124,338, September 26, 2000
- Sigma receptor ligands and the use thereof, United States Patent 6,087,346, July 11, 2000
- Benzylamidines derivatives with serotonin receptor binding activity, United States Patent 6,048,862, April 11, 2000
- Sigma receptor ligands and the use thereof. United States Patent 6,057,371, May 2, 2000
- Benzylamidines derivatives with serotonin receptor binding activity. United States Patent 5,969,137. October 19, 1999
- Methods of using pharmaceutical tetrahydroisoquinolines. United States Patent 5,919,794, July 6, 1999
- 5-HT_{1D} receptor ligands. United States Patent 5,504,101, April 2, 1996
- Tryptamine analogs with 5-HT_{1D} selectivity, United States Patent 5,496,957, March 5, 1996

2-Substituted Tryptamines: Agents with Selectivity for 5-HT₆ Serotonin Receptors^{||}

Richard A. Glennon,^{*,†} Mase Lee,[†] Jagadeesh B. Rangisetty,[†] Malgorzata Dukat,[†] Bryan L. Roth,[†] Jason E. Savage,[†] Ace McBride,[†] Laura Rauser,[†] Sandy Hufeisen,[†] and David K. H. Lee[§]

Department of Medicinal Chemistry, School of Pharmacy, Virginia Commonwealth University, Richmond, Virginia 23298, Departments of Biochemistry, Psychiatry, and Neurosciences, School of Medicine, Case Western Reserve University, Cleveland, Ohio 44106, and Allelix Biopharmaceuticals, Mississauga, Ontario L4V 1V7, Canada

Received November 4, 1999


Several 2-alkyl-5-methoxytryptamine analogues were designed and prepared as potential 5-HT₆ serotonin agonists. It was found that 5-HT₆ receptors accommodate small alkyl substituents at the indole 2-position and that the resulting compounds can bind with affinities comparable to that of serotonin. In particular, 2-ethyl-5-methoxy-*N,N*-dimethyltryptamine (**8**) binds with high affinity at human 5-HT₆ receptors ($K_i = 16$ nM) relative to 5-HT ($K_i = 75$ nM) and was a full agonist, at least as potent (**8**: $K_{act} = 3.6$ nM) as serotonin ($K_{act} = 5.0$ nM), in activating adenylate cyclase. Compound **8** displays modest affinity for several other populations of 5-HT receptors, notably h5-HT_{1A} ($K_i = 170$ nM), h5-HT_{1D} ($K_i = 290$ nM), and h5-HT₇ ($K_i = 300$ nM) receptors, but is otherwise quite selective. Compound **8** represents the first and most selective 5-HT₆ agonist reported to date. Replacing the 2-ethyl substituent with a phenyl group results in a compound that retains 5-HT₆ receptor affinity (i.e., **10**: $K_i = 20$ nM) but lacks agonist character. 2-Substituted tryptamines, then, might allow entry to a novel class of 5-HT₆ agonists and antagonists.

Introduction

Serotonin (5-hydroxytryptamine, 5-HT; **1a**) receptors are classified as belonging to one of several different families: 5-HT₁–5-HT₇. One of the newest populations identified are the 5-HT₆ receptors.¹ 5-HT₆ serotonin receptors are members of the G-protein superfamily, are positively coupled to an adenylate cyclase second-messenger system, and are found primarily in the central nervous system.² The exact clinical significance of 5-HT₆ receptors is unknown at this time. Of interest, however, is that a number of typical and atypical antipsychotic agents and tricyclic antidepressants bind with high affinity at 5-HT₆ receptors (i.e., with K_i values of <100 nM).^{3–5} In rats prevented from expressing 5-HT₆ receptors, the animals behave in a manner that seems to involve an increase in cholinergic function; this has led to speculation that one of the roles of 5-HT₆ receptors may be to control cholinergic neurotransmission and that 5-HT₆-selective antagonists could be useful in the treatment of anxiety and memory deficits.^{6,7} It has been further suggested that GABA-containing neurons in the striatum and glutamate-containing neurons in the hippocampus could be targets of 5-HT actions mediated by 5-HT₆ receptors.⁸ 5-HT₆ ligands might thus be of value in the treatment of anxiety and related disorders. Other studies suggest that 5-HT₆ receptors might be involved in motor func-

tion, mood-dependent behavior, and early growth processes involving serotonin.^{9–11}

Ro 04-6790 (**2a**) and Ro 63-0563 (**2b**) represent the first 5-HT₆-selective antagonists.¹² Several related structures have also been reported including SB-271046 (**3**, R = H).¹³ Repeated intracerebroventricular administration of antisense oligonucleotides to rats to prevent expression of 5-HT₆ receptors produces a behavioral syndrome that consists of yawning, stretching, and chewing.^{6,7} administration of Ro 04-6790 and Ro 63-0563 to naive animals produced a similar effect.¹² [³H]Ro 63-0563 has been developed as a radioligand for binding studies.^{12b}



No 5-HT₆-selective agonists have yet been identified. Various indolealkylamines, including the tryptamines


^{||} Presented in part at the Virginia Academy of Science Meeting, Norfolk, VA, May 25–28, 1999.

* Correspondence to: Richard A. Glennon, Ph.D., Box 980540, Department of Medicinal Chemistry, Virginia Commonwealth University, Richmond, VA 23298-0540. Phone: 804-828-8487. Fax: 804-828-7404. E-mail: glennon@hsc.vcu.edu.

[†] Virginia Commonwealth University.

[‡] Case Western Reserve University.

[§] Allelix Biopharmaceuticals.

Scheme 2^a

^a (a) NaH, DMF, rt, and Me₂SO₄, EtBr, nPrCl or iPrCl; (b) NaH, DMF, PhSO₂Cl, rt; (c) BuLi, DME, nPrI, -10 °C; (d) Mg, MeOH, rt; (e) γ -butyrolactone; (f) PPE, CHCl₃, reflux; (g) B₂H₆/THF, rt.

Table 1. Physicochemical Properties and 5-HT₆ Receptor Affinities of Tryptamine Analogues

compd	R	R ₃	R ₂	R ₁	yield (%)	RS ^a	mp (°C)	5-HT ₆ affinity ^b K _i (nM)	empirical formula ^c
4	H	OH	CH ₃	H				46 ^e	
5	H	OCH ₃	CH ₃	H				98	
6	CH ₃	H	CH ₃	H	88	EtOH-Et ₂ O	208	300	C ₁₄ H ₁₈ N ₂ O·HCl
7	CH ₃	OCH ₃	CH ₃	H	92	A	242–245 dec	80	C ₁₄ H ₁₈ N ₂ O·C ₂ H ₅ O ₄
8 (EMDT)	CH ₃	OCH ₃	C ₂ H ₅	H	16	B-Et ₂ O	123	52	C ₁₆ H ₂₄ N ₂ O·C ₂ H ₅ O ₄
9	CH ₃	OCH ₃	nC ₃ H ₇	H	45	A	146–147	185	C ₁₈ H ₂₄ N ₂ O·C ₂ H ₅ O ₄ ^d
10	CH ₃	OCH ₃	C ₆ H ₁₆	H	25	A	187–188	54	C ₁₈ H ₂₈ N ₂ O·C ₂ H ₅ O ₄
11	CH ₃	OCH ₃	H	H				78	
12	CH ₃	OCH ₃	H	CH ₃	65	A	181–182	510	C ₁₄ H ₂₀ N ₂ O·C ₂ H ₅ O ₄
13	CH ₃	OCH ₃	H	C ₂ H ₅	22	A	160–161	240	C ₁₅ H ₂₂ N ₂ O·1.5C ₂ H ₅ O ₄
14	CH ₃	OCH ₃	H	nC ₃ H ₇	93	A-Et ₂ O	104	200	C ₁₆ H ₂₄ N ₂ O·C ₂ H ₅ O ₄
15	CH ₃	OCH ₃	H	iC ₃ H ₇	49	A-Et ₂ O	101	130	C ₁₆ H ₂₄ N ₂ O·C ₂ H ₅ O ₄
16	CH ₃	OCH ₃	-CH ₂ CH ₂ CH ₂ CH ₂ -		75	A	114–115	1030	C ₁₇ H ₂₄ N ₂ O·1.15C ₂ H ₅ O ₄ ^e
17					37	EtOH-Et ₂ O	224–226	168	C ₁₈ H ₂₈ N ₂ O·C ₂ H ₅ O ₄

^a Recrystallization solvents: EtOH represents absolute ethanol; Et₂O, anhydrous ether; A, acetone; B, ethyl acetate. ^b K_i values represent replicate determinations and SEM are $\pm 25\%$; for purpose of comparison, clozapine was determined to possess a K_i = 5.3 (± 0.4) nM. ^c All compounds analyzed correctly to within 0.4% of theory for C, H, and N except where noted. ^d Crystallized with 0.7 mol of H₂O. ^e Crystallized with 1 mol of H₂O. ^f K_i value previously reported;¹⁶ included for purpose of comparison.

analogue of 5-methoxytryptamine (**1b**; K_i = 88 nM),¹⁶ namely, 5-methoxy-2-methyltryptamine (**5**). Compound **5** (K_i = 98 nM; Table 1) was found to bind at 5-HT₆ receptors with an affinity comparable to that of 5-methoxytryptamine. It was also found that **5** lacks affinity for 5-HT₃ receptors (K_i > 10000 nM). Compound **5** might

be a useful 5-HT₆ ligand; however, given that **5** possesses a primary amine, its utility for future in vivo studies might be hampered by its reduced ability to penetrate the blood-brain barrier and/or due to its potential for rapid metabolism by oxidative deamination. To address these problems, we sought to prepare

Table 2. Binding Profile of Compounds 7, 8, and 10^a

receptor population	<i>K_i</i> , nM (±SEM)			
	7	8 (EMDT)	10	control (agent and <i>K_i</i>)
NET	6380 (±3190)	> 10000	> 10000	nortriptyline 6.3 ± 1.2
SERT	> 10000	> 10000	4700 (±1550)	fluoxetine 3.5 ± 0.7
h5-HT _{1A}	200 (±60)	170 (±54)	1470 (±310)	WAY 100,635 0.6 ± 1.5
h5-HT _{1D}	250 (±180)	290 (±70)	6225 (±70)	ergotamine 0.8 ± 0.6
h5-HT _{1E}	1800 (±600)	520 (±180)	> 10000	serotonin 0.5 ± 0.15
r5-HT _{2A}	> 10000	> 10000	470 (±10)	clozapine 9 ± 1
r5-HT _{2C}	4020 (±640)	1810 (±490)	675 (±180)	clozapine 23 ± 5
h5-HT _{3A}	10450 (±2195)	4620 (±650)	5160 (±930)	ergotamine 22 ± 3
h5-HT ₇	145 (±34)	300 (±60)	155 (±35)	ergotamine 9 ± 2
h5-HT ₆	60 (13)	16 (±4)	20 (±5)	clozapine 10 ± 3

^a Compounds displayed *K_i* values of > 10000 nM at the following populations of receptors: histamine, NMDA, PCP, acetylcholine, opiate, and vasopressin receptors; see Experimental Section for specific subpopulations examined. *K_i* values were > 10000 nM for compounds 7 and 8 at hD₁, rD₂, rD₃, rD₄, and hD₅ receptors and > 10000 nM for 10 at hD₁, rD₂, and rD₄ receptors; although 10 produced 70% inhibition at 10000 nM at rD₃ and hD₅ receptors, it was not further evaluated. NET and SERT represent the norepinephrine and serotonin transporters. *K_i* values for all three compounds at the dopamine transporter were > 10000 nM.

several related derivatives that were somewhat more lipophilic and/or that might be less prone to metabolism.

One approach to enhancing lipophilicity and hindering metabolism was to add *N,N*-dimethyl substituents to the terminal amine; a second approach to enhancing lipophilicity was to homologate the 2-position substituent. 2-Methyl-*N,N*-dimethyltryptamine (2-methyl DMT, **6**; *K_i* = 300 nM), an *N,N*-dimethyl analogue of **5** lacking the 5-methoxy group, binds with severalfold lower affinity than **5** itself. Reintroduction of the 5-methoxy group, affording 2-methyl-5-methoxy DMT (**7**; *K_i* = 80 nM), enhanced affinity. Homologation of the 2-methyl substituent to an ethyl group (i.e., **8**; *K_i* = 52 nM) resulted in a slight increase in affinity and in a compound with affinity at least comparable to that of 5-HT itself. Further homologation of the ethyl substituent to a 2-*n*-propyl group (i.e., **9**; *K_i* = 185 nM) reversed this trend. To explore the possibility of bulk tolerance, we examined the 2-phenyl derivative **10** (*K_i* = 54 nM) and found it to bind with an affinity comparable to that of **8**.

Another attempt to enhance lipophilicity was to incorporate small alkyl substituents at the indole N₁-position. The idea was to subsequently incorporate a 2-alkyl substituent into whatever N₁-substituted analogue retained high 5-HT₆ receptor affinity. N₁-Methylation of 5-methoxy DMT (**11**; *K_i* = 78 nM) decreased 5-HT₆ receptor affinity of the resulting compound by >6-fold (**12**; *K_i* = 510 nM). Homologation of the *N*-methyl group to an ethyl group (i.e., **13**; *K_i* = 240 nM) or *n*-propyl group (i.e., **14**; *K_i* = 200 nM) doubled affinity, but the compounds did not bind as well as **11**. Branching of **13** to the isopropyl derivative **15** (*K_i* = 130 nM) resulted in a further slight enhancement of affinity. However, none of these compounds displayed significantly enhanced affinity. Compound **16** (*K_i* = 1030 nM), which may be viewed as a cyclic 1,2-disubstituted analogue of **11**, was also prepared for evaluation and was found to bind with reduced affinity.


In a final attempt to enhance lipophilicity in the 2-substituted DMT series, the propyl group of **9** was tethered to the DMT side chain to afford **17**; compound **17** (*K_i* = 168 nM) was found to bind at 5-HT₆ receptors with about 3-fold lower affinity than **8**. Although compound **17** possesses an asymmetric center and can exist as a pair of optical isomers, no attempt was made to examine the individual isomers because structurally related agents have been shown to bind at 5-HT_{1D} receptors,^{21,22} and it was anticipated that the isomers of **17** might lack the desired selectivity.

Binding Profile. Compounds **7**, **8**, and **10** were selected for examination of detailed binding profiles. All three agents were examined at more than 30 different receptor populations and produced <50% inhibition of binding at a concentration of 10000 nM at most of these populations. Where >50% displacement was observed, *K_i* values were determined (Table 2). For these studies, *K_i* values were redetermined for **7**, **8**, and **10**. Compounds **8** and **10** bind at human 5-HT₆ receptors with comparable affinity (*K_i* = 16 and 20 nM, respectively) and with an affinity similar to that of clozapine; compound **7** binds with severalfold lower affinity (*K_i* = 60 nM). Although **7** and **8** appear relatively selective, they also bind at h5-HT_{1A}, h5-HT_{1D}, h5-HT_{1E}, and h5-HT₇ receptors, yet compound **8**, in particular, still displays 10-fold selectivity over 5-HT_{1A} receptors and nearly 20-fold selectivity over h5-HT_{1D} and h5-HT₇ receptors. Compound **10** is more selective and displays 735-fold selectivity over h5-HT_{1A} receptors and >300-fold selectivity over h5-HT_{1D} receptors.

Functional Studies. Compounds **7**, **8**, and **10** were examined for their ability to activate adenylate cyclase. Whereas compounds **7** and **8** behaved as full agonists (*K_{act}* = 7.9 ± 5.0 and 3.6 ± 1.3 nM, respectively) relative to 5-HT (*K_{act}* = 5.0 ± 3.0 nM), compound **10** showed no agonist activity (see Figure 1). Compound **10** inhibited 5-HT-stimulated adenylate cyclase at 10000 nM suggesting that it is an antagonist.

Summary

Molecular manipulation of a tryptamine template revealed that 2-methyl substitution was tolerated by 5-HT₆ receptors.¹⁶ Because 2-methyl-5-HT was previously considered to be a 5-HT₃-selective ligand and by taking advantage of the fact that 5-HT₃ receptors do not readily accommodate a 5-methoxy group, a series of






Figure 1. Typical dose-response curves for the effects of compounds **7**, **8**, and **10** as 5-HT_{2a} agonists in an adenylate cyclase assay; serotonin was used as control. Each compound was examined at five concentrations.

2-alkyl-5-methoxytryptamines was synthesized for evaluation at 5-HT_{2a} receptors. Several compounds were identified with affinities at least comparable to that of 5-HT itself ($K_i = 75$ nM). In particular, 2-ethyl-5-methoxy-*N,N*-dimethyltryptamine (EMDT; **8**) possessed high affinity ($K_i = 16$ nM) and displayed reasonable selectivity for 5-HT_{2a} versus other receptors examined. In functional studies, EMDT (**8**) was demonstrated to behave as a 5-HT_{2a} agonist ($K_{act} = 3.6$ nM) with a potency at least equivalent to that of 5-HT ($K_{act} = 5.0$ nM). EMDT is the most selective 5-HT_{2a} agonist reported to date. Also of interest is the 2-phenyl derivative **10** (MPDT; $K_i = 20$ nM), which possesses a somewhat different binding profile than **8**; compound **10** lacks agonist activity up to concentrations of 10000 nM and may represent a novel 5-HT_{2a} antagonist. Indeed, when examined at the single concentration of 10000 nM, **10** behaved an antagonist. Hence, with the appropriate substituents, 2-substituted tryptamines may provide entry to new 5-HT_{2a}-selective agonists and antagonists.

Experimental Section

Synthesis. Melting points, determined with a Thomas-Hoover melting point apparatus, are uncorrected. Proton magnetic resonance spectra were obtained with a GE QE-300 or Varian Gemini 300 spectrometer; and tetramethylsilane was used as an internal standard. Infrared spectra were recorded on a Nicolet 52DX FT-IR. Elemental analysis was performed by Atlantic Microlab Inc. and determined values are within 0.4% of theory. Flash chromatography was performed on silica gel (Merck grade 60, 230–400 mesh 60 Å). Certain compounds were previously reported in the literature but due to difficulty in either preparing or purifying the reported salt, a different salt was prepared. Specifically, compounds **7**,¹⁷ **10**,²³ and **12**²⁴ are known as their HCl salts but were isolated as their monooxalate salts in the present investigation. Compound **6**, prepared earlier as a maleate salt,²⁵ was isolated as its HCl salt. All four of these compounds analyzed correctly for C, H, and N.

2-Ethyl-5-methoxy-*N,N*-dimethyltryptamine Maleate (8**).** A 2.5 M solution of *n*-BuLi (1.75 mL, 4.38 mmol) was added in a dropwise manner to a stirred solution of **7**¹⁷ (free base) (1.00 g, 4.33 mmol) in dry THF (7 mL) at -78°C under N₂. After stirring the reaction mixture for 5 min, the cooling bath was removed and CO₂ gas was passed into the solution for 10 min. The solvent was removed at 0°C under reduced pressure

to give a transparent solid. The flask was flushed with N₂ and dry THF (7 mL) was added. The reaction mixture was degassed at -150°C under reduced pressure of 1 mmHg, then allowed to warm to -78°C ; 1.7 M *n*-BuLi (2.8 mL, 4.8 mmol) was added to give a bright yellow solution. The cooling bath was replaced by an ice-salt bath and the reaction was kept at -20°C for 45 min, then cooled to -78°C , and MeI (0.3 mL, 4.81 mmol) was added in a dropwise manner. The solution was kept at -78°C for 3 h. The reaction mixture was acidified with a saturated ethereal solution of HCl. Anhydrous Et₂O was added to the resulting suspension and the supernatant was decanted. The residue was heated at 100°C under reduced pressure for 20 min. The resulting residue was purified by flash chromatography on silica gel (CH₂Cl₂/MeOH; 12:1) to give 0.17 g of a bright yellow oil (16%): ¹H NMR (CDCl₃) δ 8.06 (s, 1H), 7.14 (d, 1H, $J = 8.67$ Hz), 6.98 (s, 1H), 6.76 (dd, 1H, $J = 2.34$, 8.73 Hz), 3.84 (s, 3H), 2.91–2.87 (m, 2H), 2.71 (q, 2H, $J = 7.38$ Hz), 2.57–2.52 (m, 2H), 2.38 (s, 6H), 1.25 (t, 3H, $J = 7.38$ Hz). The maleate salt was prepared and recrystallized from an EtOAc/Et₂O mixture: mp 123°C . Anal. (C₁₅H₂₂N₂O₂·C₄H₄O₄) C, H, N.

5-Methoxy-2-*n*-propyl-*N,N*-dimethyltryptamine Oxalate (9**).** Magnesium turnings (840 mg) and NH₄Cl (77 mg, 1.44 mmol) were added to a solution of **19** (free base) (259 mg, 0.65 mmol) in MeOH (17 mL) and the mixture was allowed to stir at room temperature for 1 h. Saturated NH₄Cl solution was added and the reaction mixture was extracted with CH₂Cl₂. The organic portion was dried (MgSO₄) and the solvent was removed under reduced pressure. The residue was purified by flash chromatography on silica gel (CH₂Cl₂/MeOH; 9:1) to give 75 mg (45%) of a bright yellow oil: ¹H NMR (CDCl₃) δ 7.71 (bs, 1H), 7.16 (d, 1H, $J = 8.67$ Hz), 6.99 (d, 1H, $J = 2.43$, 8.77 Hz), 6.77 (dd, 1H, $J = 2.25$, 8.73 Hz), 3.85 (s, 3H), 2.89–2.83 (m, 2H), 2.69 (t, 2H, $J = 7.56$ Hz), 2.53–2.47 (m, 2H), 2.36 (s, 6H), 1.68 (q, 2 H, $J = 7.28$, 7.56 Hz), 0.98 (t, 3H, $J = 7.28$ Hz). The oxalate salt was prepared and recrystallized from acetone: mp 146 – 147°C . Anal. (C₁₈H₂₆N₂O₂·C₂H₂O₄·0.7H₂O) C, H, N.

5-Methoxy-2-phenyl-*N,N*-dimethyltryptamine Oxalate (10**).** 5-Methoxy-2-phenylindole²⁶ (3 g, 13.44 mmol) was added to a stirred ice-cooled solution of 1-dimethylamino-2-nitroethylene (1.56, 13.44 mmol) in trifluoroacetic acid (8 mL). The resulting mixture was allowed to stir under N₂ at room temperature for 30 min and was then poured into ice/water. The solution was extracted with EtOAc and the organic portion was washed consecutively with saturated NaHCO₃ solution, H₂O, then brine. The organic portion was dried (MgSO₄) and solvent was removed under reduced pressure. The residue was recrystallized from CH₂Cl₂/hexane to give 2.36 g (60%) of **22** as a red powder: ¹H NMR (acetone-*d*₆) δ 8.82 (bs, 1H), 8.32 (d, 1H, $J = 13.44$ Hz), 7.94 (d, 1H, $J = 13.35$ Hz), 7.69–7.41 (m, 7H), 6.98–6.94 (m, 1H), 3.92 (s, 3H); IR (KBr) 1606, 1475, 1251 cm⁻¹. A solution of **22** (2.00 g, 6.75 mmol) in dry THF (20 mL) was added in a dropwise manner to a cooled (0°C) suspension of LiAlH₄ (1.54 g, 40.5 mmol) in dry THF (40 mL) under N₂. The reaction mixture was heated at reflux for 1 h and then allowed to stand at room temperature overnight. The resulting mixture was quenched with H₂O then 15% NaOH solution. Celite was added and the solution was filtered. The solvent was removed under reduced pressure. The residue was purified by flash chromatography on silica gel (CH₂Cl₂/MeOH; 9:1) to give 1.00 g (55%) of the primary amine **23**²¹ as an oil: ¹H NMR (CDCl₃) δ 8.19 (bs, 1H), 7.59–7.58 (m, 2H), 7.49–7.44 (m, 2H), 7.39–7.34 (m, 1H), 7.28–7.25 (m, 1H), 7.09 (d, 1H, $J = 2.37$ Hz), 6.88 (dd, 1H, $J = 2.24$, 8.75 Hz), 3.89 (s, 3H), 3.04 (bs, 4H); IR (KBr) 3397, 3347 cm⁻¹. Sodium cyanoborohydride (510 mg, 8.12 mmol) was added to a solution of primary amine **23** (700 mg, 2.63 mmol) and 37% aqueous CH₂O in MeCN (10 mL) at room temperature. The resulting mixture was adjusted to pH 5 with HOAc and was allowed to stir at room temperature overnight. A 15% solution of NaOH was added to neutralize the mixture and the mixture was extracted with CH₂Cl₂. The combined organic portion was washed with saturated NaHCO₃ solution and brine. The

organic portion was dried (MgSO_4) and solvent was removed under reduced pressure. The residue was purified by flash chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{MeOH}$: 9:1) to give 195 mg (25%) of **10** (free base) as a white powder: ^1H NMR (CDCl_3) δ 8.05 (bs, 1H), 7.56–7.53 (m, 2H), 7.49–7.44 (m, 2H), 7.39–7.34 (m, 1H), 7.29–7.25 (m, 1H), 7.11 (d, 1H, J = 2.25 Hz), 6.87 (dd, 1H, J = 2.52, 8.73 Hz), 3.89 (s, 3H), 3.13–3.08 (m, 2H), 2.72–2.66 (m, 2H), 2.39 (s, 6H). Although the HCl salt has been previously reported,³ difficulties in its purification led to isolation of the product as its oxalate salt: mp 187–188 °C after recrystallization from acetone. Anal. ($\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}\cdot\text{C}_2\text{H}_2\text{O}_4$) C, H, N.

5-Methoxy-1-(2-propyl)-*N,N*-dimethyltryptamine Maleate (15). A mixture of 5-methoxy-*N,N*-dimethyltryptamine (**11**; free base) (500 mg, 2.29 mmol) and 60% NaOH (100 mg, 2.52 mmol) was heated at 100 °C under N_2 until evolution of H_2 gas ceased. The resultant mass was dissolved in anhydrous DMF (3 mL) and 2-bromopropane (0.25 mL, 2.84 mmol) was added to the solution at 0 °C. The reaction mixture was allowed to stir at room temperature for 3 h. Brine was added and the reaction mixture was extracted with CH_2Cl_2 . The organic portion was dried (MgSO_4) and the solvent was removed under reduced pressure. The residue was purified by flash chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{MeOH}$: 10:1) to give 294 mg of a bright yellow oil (49%): ^1H NMR (CDCl_3) δ 7.23 (d, 1H, J = 8.94 Hz), 7.04 (d, 1H, J = 2.46 Hz), 7.01 (s, 1H), 6.86 (dd, 1H, J = 2.46, 8.88 Hz), 4.59–4.54 (m, 1H), 3.86 (s, 3H), 2.95–2.89 (m, 2H), 2.66–2.61 (m, 2H), 2.36 (s, 6H), 1.48 (d, 6H, J = 6.72 Hz). The maleate salt was prepared and recrystallized from an acetone/ Et_2O mixture: mp 101–102 °C. Anal. ($\text{C}_{19}\text{H}_{24}\text{N}_2\text{O}\cdot\text{C}_4\text{H}_4\text{O}_4$) C, H, N.

^1H NMR data for compounds **13** and **14** are as follows: **13** (CDCl_3) δ 7.22 (d, 1H, J = 9.0 Hz), 7.06 (d, 1H, J = 2.5 Hz), 6.95 (s, 1H), 6.88 (dd, 1H, J = 2.5, 6.0 Hz), 4.10 (q, 2H, J = 7.5 Hz), 3.88 (s, 3H), 3.01–2.97 (m, 2H), 2.76–2.73 (m, 2H), 2.45 (s, 6H), 1.44 (t, 3H, J = 7.5 Hz); **14** (CDCl_3) δ 7.19 (d, 1H, J = 8.85 Hz), 7.04 (d, 1H, J = 2.37 Hz), 6.90 (s, 1H), 6.87–6.83 (m, 1H), 3.98 (t, 2H, J = 7.08 Hz), 3.86 (s, 3H), 2.93–2.87 (m, 2H), 2.64–2.59 (m, 2H), 2.35 (s, 6H), 1.82 (q, 3H, J = 7.2 Hz), 0.91 (t, 2H, J = 7.4 Hz).

6,7,8,9-Tetrahydro-2-methoxy-10-(*N,N*-dimethylamino)ethylpyrido[1,2-*a*]indole Oxalate (16). A solution of 1.0 M borane/THF (2 mL, 2 mmol) was added in a dropwise manner to ice-bath cooled **21** (290 mg, 1.01 mmol) under N_2 . The reaction mixture was allowed to stir at room temperature for 2 h. Acetone (3 mL) was added, and the reaction mixture was heated at reflux for 1 h to quench the unreacted borane reagent. The solvent was removed under reduced pressure. A 15% solution of NaOH was added and the mixture was extracted with CH_2Cl_2 , and the CH_2Cl_2 portion was washed with H_2O , then brine. Solvent was removed under reduced pressure. The residue was purified by flash chromatography on silica gel (hexane/ EtOAc : 4:1) to give 207 mg (75%) of a light yellow oil: ^1H NMR ($\text{DMSO}-d_6$) δ 7.34 (d, 1H, J = 8.85 Hz), 7.21 (s, 1H), 7.11 (s, 1H), 4.08 (t, 2H, J = 6.65 Hz), 3.79 (s, 3H), 3.40–3.35 (m, 2H), 3.30–3.25 (m, 2H), 3.06–3.01 (m, 2H), 2.83 (s, 6H), 1.76–1.69 (m, 2H), 1.40–1.31 (m, 2H). A small portion was converted to its oxalate salt: mp 114–115 °C. Anal. ($\text{C}_{27}\text{H}_{28}\text{N}_4\text{O}\cdot 1.5\text{C}_2\text{H}_2\text{O}_4\cdot\text{H}_2\text{O}$) C, H, N.

4-(Dimethylamino)ethyl-6-methoxy-1,2,3,4-tetrahydrocarbazole Oxalate (17). Sodium metal (1.0 g) was added portionwise over a 30-min period to a stirred solution of 4-(dimethylamino)ethyl-9-benzyl-6-methoxy-1,2,3,4-tetrahydrocarbazole hydrochloride¹⁹ (4.0 g, 0.01 mol) in liquid NH_3 (300 mL). NH_4Cl (3.0 g) was added until the blue color of the mixture disappeared. The NH_3 was evaporated, H_2O (50 mL) was added, and the mixture was extracted with CH_2Cl_2 (3 \times 50 mL). The combined organic portion was washed with H_2O (50 mL), brine (50 mL), dried (MgSO_4) and evaporated to give an oil. The oil was purified by column chromatography ($\text{CHCl}_3/\text{MeOH}$: 9:1) and converted to an oxalate salt. The oxalate salt was recrystallized from anhydrous Et_2O /absolute EtOH to give 1.8 g (37%) of the desired target as a white powder: mp 224–226 °C; ^1H NMR (CDCl_3 , free base) δ 8.10 (s, 1H, NH), 7.20 (t,

1H, ArH), 6.90 (d, 1H, ArH), 6.70 (dd, 1H, ArH), 3.80 (s, 3H, OCH_3), 3.40 (t, 1H, CH), 3.15 (d, 1H, CH), 3.00 (t, 1H, CH), 2.82 (s, 6H, $2 \times \text{CH}_3$), 2.63–2.73 (m, 2H, CH), 2.33 (m, 1H, CH), 1.8–2.0 (m, 3H, $\text{CH}_2\text{-CH}$). Anal. ($\text{C}_{16}\text{H}_{22}\text{N}_2\text{O}\cdot\text{C}_2\text{H}_2\text{O}_4$) C, H, N.

1-Benzenesulfonyl-5-methoxy-*N,N*-dimethyltryptamine Oxalate (18). A mixture of 5-methoxy-*N,N*-dimethyltryptamine (**11**; free base) (4.35 g, 19.93 mmol) and 60% NaOH (0.87 g, 21.75 mmol) was heated at 100 °C under N_2 until evolution of H_2 gas ceased. The resultant mass was dissolved in anhydrous DMF (21 mL) and benzenesulfonyl chloride (2.8 mL, 21.94 mmol) was added in a dropwise manner at 0 °C. The reaction mixture was allowed to stir at room temperature overnight. Saturated NaHCO_3 solution was added and the mixture was extracted with CH_2Cl_2 . The organic portion was dried (MgSO_4) and the solvent was removed under reduced pressure. The residue was purified by flash chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{MeOH}$: 9:1) to give 4.39 g of an oil (61%): ^1H NMR (CDCl_3) δ 7.89–7.87 (m, 1H), 7.83 (d, 2H, J = 8.0 Hz), 7.51 (t, 1H, J = 7.8 Hz), 7.34 (s, 1H), 6.93–6.92 (m, 2H), 3.82 (s, 3H), 2.80 (t, 2H, J = 7.8 Hz), 2.59 (t, 2H, J = 7.8 Hz), 2.33 (s, 6H); IR (CHCl_3) 1357, 1115 cm^{-1} . The oxalate salt was prepared and recrystallized from an acetone/ Et_2O mixture: mp 224–226 °C. Anal. ($\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_5\cdot\text{C}_2\text{H}_2\text{O}_4$) C, H, N.

1-Benzenesulfonyl-5-methoxy-2-*n*-propyl-*N,N*-dimethyltryptamine Oxalate (19). A 2.5 M solution of *n*-BuLi (1.4 mL, 3.5 mmol) was added in a dropwise manner to a stirred solution of **18** (free base) (1.00 g, 2.79 mmol) in DMF (4 mL) at -10 °C under N_2 . The resulting solution was allowed to stir for an additional 10 min at -10 °C, and then *n*-PrI (0.35 mL, 3.59 mmol) was added. The reaction mixture was allowed to stir for 1 h at -10 °C. Saturated NaHCO_3 solution was added and the reaction mixture was extracted with CH_2Cl_2 . The organic portion was washed with brine and dried (MgSO_4); the solvent was removed under reduced pressure and the residue was purified by flash chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{MeOH}$: 30:1) to give 0.19 g (17%) of a bright yellow oil: ^1H NMR (CDCl_3) δ 8.06 (d, 1H, J = 8.79 Hz), 7.62 (d, 2H, J = 8.22 Hz), 7.51–7.46 (m, 1H), 7.38–7.33 (m, 2H), 6.95 (bs, 1H), 6.89–6.85 (m, 1H), 3.85 (s, 3H), 2.96–2.89 (m, 4H), 2.63–2.57 (m, 2H), 2.48 (s, 6H), 1.73 (q, 2H, J = 7.51 Hz), 1.00 (t, 3H, J = 7.51 Hz); IR (CHCl_3) 1355 cm^{-1} . The oxalate salt was prepared and recrystallized from acetone: mp 175–176 °C. Anal. ($\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_5\cdot\text{C}_2\text{H}_2\text{O}_4$) C, H, N.

6,7,8,9-Tetrahydro-2-methoxy-10-(*N,N*-dimethylamino)ethylpyrido[1,2-*a*]indol-9-one Oxalate (21). A mixture of 5-methoxy-*N,N*-dimethyltryptamine (**11**; free base) (2.00 g, 9.17 mmol) and 60% NaOH (0.41 g, 10.1 mmol) was heated at 100 °C under N_2 until evolution of H_2 gas ceased. The resultant mass was dissolved in anhydrous DMF (25 mL) and anhydrous γ -butyrolactone (1.4 mL, 18.2 mmol) was added in a dropwise manner at room temperature. The reaction mixture was heated at reflux for 20 h, cooled to 0 °C, and acidified by the addition of a saturated ethereal solution of HCl. Additional Et_2O was added to the resulting suspension and the supernatant was decanted. The residue was dissolved in PPE (52.5 mL) and CHCl_3 (100 mL) and the reaction mixture was heated at reflux for 3 h under N_2 . The resulting mixture was neutralized by the addition of 15% NaOH solution, at ice-bath temperature, and extracted with CH_2Cl_2 . The organic portion was dried (MgSO_4) and solvent was removed under reduced pressure. The residue was purified by flash chromatography on silica gel ($\text{CH}_2\text{Cl}_2/\text{MeOH}$: 20:1) to give 0.52 g (20%) of **21** (free base) as a yellow oil: ^1H NMR ($\text{DMSO}-d_6$) δ 7.35 (d, 1H, J = 8.79 Hz), 7.18 (s, 1H), 6.88 (d, 1H, J = 8.85 Hz), 4.06 (t, 2H, J = 6.60 Hz), 3.80 (s, 3H), 3.42–3.36 (m, 2H), 3.17–3.12 (m, 2H), 2.85 (s, 6H), 2.66–2.62 (m, 2H); IR (CHCl_3) 1648 cm^{-1} . A small sample was converted to the oxalate salt: mp 191–192 °C dec. Anal. ($\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_5\cdot 1.6\text{C}_2\text{H}_2\text{O}_4$) C, H, N.

5-HT₆ Radioligand Binding Assay. The binding assay employed human 5-HT₆ receptors stably transfected to HEK 293 human embryonic kidney cells with PHLYsergic acid diethylamide (70 Ci/mmol; DuPont NEN) as radioligand. All

assays were conducted in triplicate using polypropylene 1 mL/well plates (Anachem). The radioligand was diluted in incubation buffer in borosilicate glass vials and protected from light. Competing agents (1 mM stock solutions) were dissolved in DMSO or saline and stored at -20°C in 1.2-mL polypropylene tubes (Eikay). Dilutions of compounds were made using incubation buffer in 96-well polypropylene plates and mixed by multichannel pipetting >25 times. Serial dilutions (1 in 4) started at a final concentration of 10000 nM. Final concentrations >10000 nM were individually prepared from the 1 mM stock solution. Nonspecific binding was defined by 100 nM serotonin creatinine sulfate (Research Biochemicals) prepared fresh in incubation buffer at the time of each determination and protected from light. Reactions volumes were as follows: 200 μL of incubation buffer (50 mM Tris, 0.5 mM EDTA, 10 mM MgCl_2), pH 7.4 at 22°C , 100 μL of test agent or serotonin (100 μM) or buffer (for total binding), 100 μL of [^3H]lysergic acid diethylamide (2 nM final concentration), and 100 μL of membrane preparation (15 μg of protein). The incubation was initiated by the addition of membrane homogenate and the plates were vortexed (Baxter S/P multibute mixer). The plates were incubated with protection from light, by shaking (Gyrotop water bath/shaker model C76, speed 2) at 37°C for 60 min. The binding reaction was stopped by filtration. The samples were filtered under vacuum over 96-well glass fiber filters (Packard Unifilter GF/B), presoaked in 0.3% PEI in 50 mM Tris buffer (4 $^{\circ}\text{C}$, pH 7.4) for at least 1 h, and then washed six times with 1 mL of cold 50 mM Tris buffer (pH 7.4) using the Packard Filtermate 96 harvester. The Unifilter plates were dried overnight in a 37°C dry incubator. The Unifilter bottoms were sealed and 35 μL of Packard MicroScint-0 was added. The plates were allowed to equilibrate for 1 h and were then sealed using a Packard TopSeal P with the Packard Plate Micromate 496. Plates were counted in a Packard TopCount 4.1 by liquid scintillation spectrometry. Each well was counted for 3 min. The test agents were initially assayed at 1000 and 100 nM. If the compound was active (defined as causing at least 80% inhibition of [^3H]lysergic acid diethylamide binding at 1000 nM), it was further tested for determination of a K_i value. The range of concentrations was chosen such that the middle concentration would produce approximately 50% inhibition.

Receptor Screen. Assays for the following receptors were performed by the NIMH Psychoactive Drug Screening Program: (1) serotonin receptors: 5-HT_{1A} , 5-HT_{1B} , 5-HT_{1D} , 5-HT_{1E} , 5-HT_{2A} , 5-HT_{2C} , 5-HT_{2E} , 5-HT_{2F} ; (2) dopamine receptors: hD_1 , rD_2 , rD_3 , rD_4 , hD_5 ; (3) muscarinic acetylcholine receptors: hM_1 , hM_2 , hM_3 , hM_4 , hM_5 ; (4) nicotinic acetylcholine receptors: $\alpha 2/\beta 2$, $\alpha 2/\beta 4$, $\alpha 3/\beta 2$, $\alpha 3/\beta 4$, $\alpha 4/\beta 2$, $\alpha 4/\beta 4$; (5) vasopressin receptors: hV_1 , hV_2 , hV_3 ; (6) opiate receptors: $\text{h}\mu$, $\text{h}\delta$, $\text{r}\kappa$; (7) transporters: hSERT, hNET, rDAT; (8) nNMDA; (9) rPCP; and (10) rH1-histamine. Detailed on-line protocols for the binding assays are described at: <http://meds20785.cwr.edu/myweb/protocol.htm>. For screening purposes, the ability of 10 μM of each compound (dissolved in 10% DMSO) was incubated with the appropriate receptor preparation and percent inhibition determined for duplicate determinations each performed in duplicate. Where $>50\%$ inhibition of specific binding was measured, K_i determinations were then measured by competition binding assays in which concentrations from 1 to 10000 nM were incubated in duplicate. For each K_i value the data represent the mean \pm SD of computer-derived estimates for $N = 4$ separate determinations. 5-HT_A receptor assays were performed exactly as previously described.³⁴

Adenylyl Cyclase Assay. 5-HT_A receptors stably expressed in HEK-293 cells were grown in 24-well plates to near-confluency and 18 h prior to assay the medium was replaced with DMEM containing dialyzed 10% fetal calf serum. For the assay, the medium was aspirated and replaced with fresh DMEM without serum and incubated with various concentrations of test agent in a total volume of 0.5 mL for 15 min. The assay was terminated by aspiration and the addition of 10% trichloroacetic acid (TCA). The TCA extract was used for cAMP

determinations. Data represent the mean of $N = 4$ separate determinations.

Acknowledgment. Supported in part by MH57635 and MH01366 (B.L.R.) and the NIMH Psychoactive Drug Screening Program.

References

- (1) Ruat, M.; Traiffort, E.; Arrang, J.-M.; Tardivel-Lacombe, J.; Diaz, L.; Leurs, R.; Schwartz, J.-C. A novel rat serotonin (5-HT_A) receptor: molecular cloning, localization, and stimulation of cAMP accumulation. *Biochim. Biophys. Acta* **1993**, *193*, 268–276.
- (2) Sleight, A. J.; Boess, F. G.; Bos, M.; Levett-Traiffort, B.; Boursan, A. The 5-hydroxytryptamine receptors: localization and function. *Exp. Opin. Ther. Patents* **1998**, *8*, 1217–1224.
- (3) Cohen, R.; Metcalf, M. A.; Khan, N.; Druck, T.; Huebner, K.; Lachowicz, J. E.; Meltzer, H. Y.; Sibley, D. R.; Roth, B. H.; Hamblin, M. W. Cloning, characterization and chromosomal localization of a human 5-HT_A serotonin receptor. *J. Neurochem.* **1998**, *66*, 47–56.
- (4) Roth, B. L.; Craig, S. C.; Choudhary, M. S.; Uluer, A.; Monsma, F. J.; Shen, Y.; Meltzer, H. Y.; Sibley, D. R. Binding of typical and atypical antipsychotic agents to 5-hydroxytryptamine-6 and 5-hydroxytryptamine-7 receptors. *J. Pharmacol. Exp. Ther.* **1994**, *268*, 1403–1410.
- (5) Grant, G. E.; Snowman, A.; Sibley, D. R.; Snyder, S. H. Clozapine: selective labeling of sites resembling 5-HT_A serotonin receptors may reflect psychoactive profile. *Mol. Med.* **1995**, *1*, 398–406.
- (6) Boursan, A.; Borroni, E.; Austin, R. H.; Monsma, F. J.; Sleight, A. J. Determination of the role of 5-HT_A receptors in the rat brain: A study using antisense oligonucleotides. *J. Pharmacol. Exp. Ther.* **1995**, *274*, 173–180.
- (7) Sleight, A. J.; Monsma, F. J.; Borroni, E.; Austin, R. H.; Boursan, A. Effects of altered 5-ht_A expression in the rat: functional studies using antisense oligonucleotides. *Behav. Brain Res.* **1996**, *73*, 245–248.
- (8) Hamon, M.; Doucet, E.; Lefevre, K.; Miquel, M.-C.; Lanfumey, L.; Insausti, R.; Frechilla, D.; Del Rio, J.; Verge, D. Antisense and antisense oligonucleotide for probing the distribution and putative functions of central 5-HT_A receptors. *Neuropsychopharmacology* **1999**, *21*, 685–765.
- (9) Yoshioka, M.; Matsumoto, M.; Nagashima, H.; Mori, K.; Saito, H. Central distribution and function of 5-HT_A receptor subtype in the rat brain. *Life Sci.* **1998**, *62*, 1473–1477.
- (10) Gerard, C.; Martres, M.-P.; Lefevre, K.; Miquel, M.-C.; Verge, D.; Lanfumey, L.; Doucet, E.; Hamon, M.; El Mestikawy, S. Immunolocalization of serotonin 5-HT_A receptor-like material in the rat central nervous system. *Brain Res.* **1997**, *746*, 207–219.
- (11) Grimaldi, B.; Bonnin, A.; Fillion, M.-P.; Ruat, M.; Traiffort, E.; Fillion, G. Characterization of 5-ht_A receptor mRNA in the rat brain during ontogenetic development. *Neurosci-Schmeldeberg's Arch. Pharmacol.* **1998**, *357*, 393–400.
- (12) (a) Sleight, A. J.; Boess, F. G.; Bos, M.; Levett-Traiffort, B.; Riemer, C.; Boursan, A. Characterization of Ro 04-6780 and Ro 63-0563: potent and selective antagonists at human 5-HT_A receptors. *Br. J. Pharmacol.* **1998**, *124*, 556–562. (b) Boess, F. G.; Riemer, C.; Bos, M.; Bentley, J.; Boursan, A.; Sleight, A. J. The 5-hydroxytryptamine receptor-selective radioligand [^3H]Ro 63-0563 labels 5-hydroxytryptamine receptor binding sites in rat and porcine striatum. *Mol. Pharmacol.* **1998**, *54*, 577–583.
- (13) Bromidge, S. M.; Brown, A. M.; Clarke, S. E.; Dodgson, K.; Gager, T.; Grassam, H. L.; Jeffrey, P. M.; Joiner, G. F.; King, F. D.; Middleton, D. N.; Mess, S. F.; Newan, H.; Riley, G.; Routledge, C.; Wyman, P. 5-Chloro- N -(4-methoxy-3-piperazin-1-ylphenyl)-3-methyl-2-benzothienophenylamide (SB-210461): A potent, selective, and orally bioavailable 5-HT_A receptor antagonist. *J. Med. Chem.* **1999**, *42*, 202–205.
- (14) Monsma, F. J.; Shetty, Y.; Ward, R. P.; Hamblin, M. W.; Sibley, D. R. Cloning and expression of a novel serotonin receptor with high affinity for tricyclic psychotropic drugs. *Mol. Pharmacol.* **1993**, *43*, 320–327.
- (15) Glennon, R. A.; Dukak, M.; Westkaemper, R. B. Serotonin receptors and ligands. In *Psychopharmacology*; Watson, S. J., Ed.; Lippincott-Raven: New York, 1998; CD-ROM version.
- (16) Glennon, R. A.; Bondarev, M.; Roth, B. 5-HT_A serotonin receptor binding of indolealkylamines: A preliminary structure-affinity investigation. *Med. Chem. Res.* **1999**, *9*, 108–117.
- (17) Shaw, E. The synthesis of tryptamines related to serotonin. *J. Am. Chem. Soc.* **1955**, *77*, 4319–4324.
- (18) *Polychlorinated ester (PPE) in Reagents for Organic Synthesis*, Volume 1; Fieser, L. F.; Fieser, M., Eds.; John Wiley and Sons: New York, 1967; pp 892–894.

- (19) Alexander, E.; Mooradian, A. 4-Aminomethyl-9-benzyl-1,2,3,4-tetrahydrocarbazoles. United States Patent 3,939,177, Feb 17, 1976.
- (20) Gozali, H. 5-HT₃ receptors. In *Serotonin Receptors and their Ligands*; Olivier, B., van Wijngaarden, I., Soudijn, W., Eds.; Elsevier: Amsterdam, 1997; pp 221–258.
- (21) King, F. D.; Brown, A. M.; Gaster, L. M.; Kaumann, A. J.; Medhurst, A. D.; Parker, S. G.; Parson, A. A.; Patch, T. L.; Raval, P. (±)3-Amino-6-carboxamido-1,2,3,4-tetrahydrocarbazole: A conformationally restricted analogue of 5-carboxamidotryptamine with selectivity for the serotonin 5-HT₁₀ receptor. *J. Med. Chem.* 1993, **36**, 1918–1919.
- (22) Glennon, R. A.; Hong, S. S.; Bondarev, M.; Law, H.; Dukat, M.; Rakhit, S.; Power, P.; Fan, E.; Kinneau, D.; Kamboj, R.; Teitler, M.; Herrick-Davis, K.; Smith, C. Binding of O-alkyl derivatives of serotonin at human 5-HT₁₀ receptors. *J. Med. Chem.* 1996, **39**, 314–322.
- (23) Julia, M.; Melamed, R.; Gombert, R. Research in the indole series. XVI. 2-Aryltryptamines and homologous amines. *Ann. Inst. Pasteur* 1965, **109**, 343–362; *Chem. Abstr.* 1966, **64**, 677.
- (24) Benington, F.; Morin, R. D.; Clark, L. C. Jr. Synthesis of O- and N-methylated derivatives of 5-hydroxytryptamine. *J. Org. Chem.* 1958, **23**, 1977–1979.
- (25) Chapman, N. B.; Clarke, K.; Hughes, H. Synthesis of some 5-substituted 2-methyltryptamines and their N-mono- and di-alkyl derivatives. *J. Chem. Soc.* 1965, 1424–1428.
- (26) Houltham, W. J.; Parrino, V. A.; Ulke, Y. Lithiation of N-(2-alkylphenyl)alkanamides and related compounds. Madelung indole synthesis. *J. Org. Chem.* 1981, **46**, 4511–4515.

JM990550B